Case Study 1: An Evidence-Based Practice Review Report

Theme: School (setting) based interventions for children with special educational needs (SEN)

How effective is CBTp in improving psychotic symptoms and functioning in adolescents and young people below the age of 25?

Psychosis is a life-altering mental health condition that can present early in life and impact children and young people (CYP) at a time of great developmental change. Cognitive Behavioural Therapy for psychosis (CBTp) is recommended as a psychological intervention for CYP, though evidence on CBTp use in non-adult populations is scarce. During CBTp, the CYP develops a therapeutic alliance with the therapist where psychotic incidents and the beliefs the CYP holds about them are explored and ultimately replaced by more adaptive beliefs and behaviours. The goal of CBTp is to reduce the distress associated with delusions and hallucinations, with the aim of enhancing symptomatic relief and functional recovery. This review sought to explore the current literature and evaluate the effect of CBTp on psychosis and functioning in CYP under the age of 25. A systematic literature search identified five studies for review and a combination of small to large effect sizes supporting CBTp efficacy were found. Study findings and future research needs are discussed.

Cognitive Behavioural Therapy for Psychosis (CBTp) evolved from Beck's cognitive model of depression (1979) which views pathology as a product of four interacting systems: our thoughts, emotions, behaviour and physiology. CBT suggests that it is the person's interpretation of an event that affects how they feel and behave, rather than the event itself (cognitive principle), while also explaining that what a person chooses to do (behavioural principle) will in turn affect how they feel and think (Kennerley et al., 2016) (Figure 1). For example, an inaccurate belief may lead a person to avoid a certain experience, which can in turn reduce their opportunities for reality testing, thus reinforcing the original inaccurate belief, contributing to pathology.

But where does pathology originate from? The guiding theory behind both CBT and CBTp is that bio-psycho-social factors such as early childhood experiences develop our core beliefs and structures of the world, which can be functional and adaptive or maladaptive. Beck et al. (1990) explained that these maladaptive schemas develop over the years and 'await' to be triggered by critical incidents in life, in turn evoking dysfunctional behaviours or emotional states such as anxiety. On this basis, CBT ''helps people identify and change thinking and behavior patterns that are harmful or ineffective, replacing them with more accurate thoughts and functional behaviours'' (VandenBos, 2015).

Similarly, CBT models of psychosis propose that psychotic-like experiences are influenced by maladaptive appraisals of neutral or anomalous experiences, which in



CBT has been gaining recognition over the last 20 years as a potential therapy for people with psychosis (Cormac et al., 2002) and is now recommended by UK NICE guidelines (2016) as a non-pharmacological intervention for CYP experiencing psychosis. However, this recommendation is guided by the continuously growing

evidence-base supporting CBTp use in adult populations (Naeem et al., 2014) and not CBTp research in CYP, which is shockingly scarce (Stafford et al., 2015).

Early onset psychosis which is defined as the presence of a first-episode of psychosis before the age of 18 is thought to affect approximately 6 in 100,000 people in the UK (Boeing et al., 2007). Symptoms are divided into positive (e.g. delusions, hallucinations) and negative (e.g. emotional and social withdrawal, poverty of speech) (Andreasen, 1982; Passby & Broome, 2017 .3 Tbyen,nt(t)17 (i)0.9 (a)-10.]TJ 0.018 Tw (sy (i)0.9 (a

2. Adolescent Psychosis
3. Psychotic-like experiences
4. Early onset psychosis
5. Early onset schizophrenia
6. Child schizophrenia
7. Children psychosis
8. Unusual experiences
9. Child psychosis
10. Children schizophrenia
11. First episode psychosis
12. Early psychosis
13. VEOS
14. Psychosis
15. Schizophrenia
17. CBT
18. Cognitive behavioral therapy
19. Psychological intervention
20. Psychological interventions
21. Cognitive Behavioural Treatment
22. Cognitive Behavioral Treatment
23. Psychological Treatment
24. Psychological Treatments
25. Psychosocial intervention
26. Psychosocial interventions
27. Psychosocial Treatment
28. Psychosocial Treatments
29. Cognitive behaviour therapy
30. Cognitive behavior therapy

31. 1 OR 2 OR 3 OR 4 OR 5 OR 6 OR 7 OR 8 OR 9 OR 10 OR 11 OR 12 OR 13 OR 14 OR 15

32. 16 OR 17 OR 18 OR 19 OR 20 OR 21 OR 22 OR 23 OR 24 OR 25 OR 26 OR 27 OR 28 OR 29 OR 30


Identification
Screening
Eligibility
Included
Records identified through database searching (n = 1688)
Additional records identified through other sources (n = 0)
Records after duplicates removed (n= 779)
Records screened (n =779)
Records excluded (n = 653)
Full-text articles assessed for eligibility (n = 126)
Full-text articles excluded (n = 121)
Studies included in narrative synthesis
(n = 5)

1. Type of publication	The article has been published in a peer-reviewed journal	The article has not been published in a peer-reviewed journal	The article has been subjected to quality control by experts in the field
2. Study design	RCT with pre- and post-intervention measures	Non-RCT (i.e. Quasi-experimental studies, systematic reviews and meta-analyses, case experimental studies, qualitative studies	RCTs minimise the risk of confounding factors influencing the trial results and are considered to provide the most reliable evidence on the effectiveness of interventions
3. Language	Article is written in English	Article is not written in English	Author accessibility to article
4. Intervention	Trials investigating CBT-based interventions	Trials not investigating CBT-based interventions	This review seeks to evaluate the efficacy of CBT-based interventions on psychosis and functioning in those aged ≤25 years of age

Browning, S., Corrigall, R., Garety, P., Emsley, R., & Jolley, S. (2013). Psychological interventions for adolescent psychosis: A pilot controlled trial in routine care. *European Psychiatry, 28*(7), 423-426. https://doi.org/10.1016/j.eurpsy.2013.05.008

randomisation, double blinding status and their

			(N)		***(n)***			
Browning et al. (2013) U.K.	Pilot RCT	Psychotic disorder, ICD-10	CBTpA+Standard Care *(n=10)* ***Comparison Group(s)*** Family Therapy (FTpA) + inpatient Standard Care (SC). Duration: 5h long sessions over 4–10 weeks (n=10) Standard Care (SC) (n=10)	CBTpA: 10 half-hour sessions, twice a week - duration was patient length of stay in setting (M=65.5 days)	*14 - 17* *n= 30*	a) Affective and psychotic symptoms (BPRS) b) Psychosocial functioning (C-GAS)	No significant difference comparisons ( ). Improvement in symptoms and psychosocial functioning found in Significant over time in symptoms and functioning for both therapeutic groups. Post-intervention improvements only in symptoms but not functioning for the SC group.	1.60 - Medium

Muller et al. (2020) Germany	Pilot RCT	Current diagnosis of schizophrenia, schizophreniform, schizoaffective or delusional disorder (DSM-IV) according to the SCID-PD	CBTpA + Treatment as Usual (TAU) *(n= 13)* Treatment as usual (TAU) (n= 12)	9 months (20 individual sessions) of CBTpA + TAU	14-20 *n= 25*	a) PANSS-positive subscale b) PANSS - negative subscale c) Auditory Hallucinations (PSYRATS) d) Delusions (PSYRATS) e) Depression (CDSS) f) Functioning (GAF) g) Quality of life (MSQoL-R)	were trivial (d= 0.04) for the PANSS-positive subscale and lacked statistical significance. However, small between-group effect sizes in favour of CBT+TAU were seen for the PANSS negative subscale, delusions, depression, psychosocial functioning and QoL. were still trivial (d= 0.02) for the PANSS-positive subscale and lacked statistical significance. However, small between-group effect sizes in favour of CBT+TAU were seen for the “material quality of life” subscale, the PANSS negative subscale, auditory	

A total of 268 participants of which 55% were male, took part in the five studies. The median sample size was n=61 (range 25 to 90), while the participant mean age was 18.2 years (range 16.3 to 22.3). All participants shared an ICD-10 or DSM-IV diagnosis of psychosis and had active psychotic symptoms upon study entry. In their majority, participants were outpatients in specialist psychosis or adolescent services across the U.K. (Morrison et al., 2020), Australia (Francey et al., 2020; Jackson et al.,,,,,

item BPRS scale, was adopted by two of the trials (Morrison et al., 2020; Muller et al., 2020). Though the authors did not provide current or historical sources on the psychometric quality of the tools other than references to their origin, both the BPRS and PANNS are established measures, used internationally in both a clinical and research capacity, with information on their reliability and validity being readily available. Their p

Though a fairly new measure, the SPEQ has been found to have good convergent validity, test-retest reliability, and internal consistency (Ronald et al., 2014, Zavos et al., 2014). Data on the psychometric properties of the PSYRATS are also positive, with research supporting the tool for re-test and inter-rater reliability and validity, in particular, internal consistency and sensitivity to change (Drake et al., 2007; Favrod et al., 2012; Hatton et al., 2005; Mortan Sevi et al., 2016).

Francey et al. (2020) utilised the Heinrich Quality of Life Scale-QLS (Heinrichs et al., 1984) and the Social and Occupational Functioning Scale-SOFAS (APA, 1994) to measure functioning, with the latter also used to assess functioning in the Jackson et al. (2008) trial. Muller et al. (2020) assessed functioning with the Global Assessment of Functioning Scale-th the latsuncasand O ((as)u.)2 (n (unc)49 (as)(unc)4TJ [(g)2 ( )]T0 Tw (-)Tj

Study design was evaluated against WoE A and B criteria. Each study followed a randomised controlled trial design, a gold standard for investigating cause-effect relationships by stratifying participants into ‘equally’ comparable groups, while

Though this increased risk of bias for the four remaining trials and relevant points were deducted, the studies minimised bias from further influencing findings by either blinding their outcome measure raters to treatment allocation (Jackson et al., 2008; Morrison et al., 2020; Muller et al., 2020) or ensuring they were independent to CBT delivery (Browning et al., 2013; Francey et al., 2020).

All five studies provided information on their withdrawal and drop-out rates where they occurred. As missing or incomplete outcome data increase the risk of attrition bias by minimising group comparability, different methods were adopted by the studies to adjust forw 4.46 0 Td A38 Tw 1eE((udi)16 (esi43t 1y0udi)128o1 (c1f)]a.n6udI da.n6ud(e t

training, the quality of the sessions nor on treatment adherence, significantly impacting

depression, anxiety and positive and negative symptoms, while improvements were seen in their functioning and quality of life measures.

Within-group improvements in functioning and psychopathology at post-treatment and follow-up were seen in all five trials for participants receiving CBT. Given the trials received medium to high WoE D scores, relevant merit should be given to the findings and their importance to the review question. However, it is important to highlight that the trials were not powered for efficacy testing and as such their findings should be interpreted with caution.

A summary of main findings, measures and effect sizes can be found below.

N		(Cohen's d)	(Cohen, 1992)		
Browning et al. (2013) U.K. ***N*** CBTpA, n=10 ***N*** FTpA, n= 10 SC, n= 10	<u>Pre-treatment to post-treatment:</u> Affective and psychotic symptoms (BPRS) Psychosocial Functioning (C-GAS) <u>Pre-treatment to post-treatment</u> <u>(M=65.5 days):</u> Affective and psychotic symptoms (BPRS) Psychosocial Functioning (C-GAS)	Cohen's d= 0.6			

N		(Cohen's d)	(Cohen, 1992)		
Morrison et al. (2020) U.K. CBT+OFI, n , n					

N		(Cohen's d)	(Cohen, 1992)		

=Standard Care, =Cognitive Behavioural Therapy for Adolescents with Psychosis, =Family Therapy for Adolescents with Psychosis, =Brief Psychiatric Rating Scale, =Children's Global Assessment Scale, =Structured Clinical Interview for DSM-5 Personality Disorders, =Treatment As Usual, =cognitive behavioral therapy, =positive and negative syndrome scale- positive scale, = positive and negative syndrome scale-negative scale, = Psychotic Symptoms Rating Scales, =Calgary Depression Scale for Schizophrenia, =Global assessment of functioning scale, =Modular System for Quality of Life, = Optional Family Intervention = Specific Psychotic Experiences Questionnaire, = Hospital Anxiety and Depression Scale, = Active Cognitive Therapy, = Befriending =Scale for the Assessment of Negative Symptoms,

CBTp without antipsychotic medication can be effective in reducing psychotic symptoms and improve functioning in some young people with psychosis following treatment, with small to large effect sizes still evident at follow-up. Though further safety and efficacy trials replicating this data are undoubtedly needed, managing psychosis through non-pharmacological interventions would perhaps be a welcome option for many. Antipsychotic medication use in CYP is currently associated with a range of adverse effects such as weight gain (Correll et al., 2009), stigma (Murphy et al., 2015), cardiovascular risk and movement disorders (Haddad & Sharma, 2007), while their risk-benefit ratio has been found to be poor (Stafford et al., 2015). sucl.

Addington, J., & Gleeson, J. (2005). Implementing cognitive-behavioural therapy for first-episode psychosis. *The British Journal of Psychiatry, 187*(48), 72–76. https://doi.org/10.1192/bjp.187.48.s72

Alonso, J., Ciudad, A., Casado, A., & Gilaberte, I. (2008). Measuring schizophrenia remission in clinical practice. *Canadian Journal of Psychiatry*, *53*(3), 202–206. https://doi.org/10.1177/070674370805300311

American Psychiatric Association [APA]. (1994). *Diagnostic and statistical manual of* [redacted]ton, DC

Andreasen, N. C., Flaum, M., Arndt, S., Alliger, R., Swayze, V. W. (1991). *Positive and Negative Symptoms: Assessment and Validity*. In: Marneros, A., Andreasen, N. C., Tsuang, M. T. (eds) Negative Versus Positive Schizophrenia. Springer, Berlin, Heidelberg. https://doi.org/10.1007/978-3-642-76841-5_3

Ascher-Svanum, H., Montgomery, W. S., McDonnell, D. P., Coleman, K. A., & Feldman, P.D. (2012). Treatment-completion rates with olanzapine long-acting injection versus risperidone long-acting injection in a 12-month, open label treatment of schizophrenia: indirect, exploratory comparisons. *International Journal of General Medicine*, 5, 391–398

Baldwin, L. (2018). *Research Concepts for the Practitioner of Educational Leadership*. Leiden, The Netherlands: Brill. doi: https://doi.org/10.1163/9789004365155

Barker, C., Pistrang, N., & Elliott, R. R. (2005). *Research methods in clinical psychology: An introduction for students and practitioners (2nd ed.)*. Wiley-Blackwell.

Barnett, A. G., van der Pols, J. C., & Dobson, A. J. (2005). Regression to the mean: what it is and how to deal with it. *International Journal of E*

Beck, A. T., Rush, A. J., Shaw, B. F., & Emery, G. (1979).

Clemmensen, L., Vernal, D. L., & Steinhausen, H. C. (2012). A systematic review of the long-term outcome of early onset schizophrenia. *BMC Psychiatry, 12,* 150. https://doi.org/10.1186/1471-244X-12-150

Cohen, J. (1988), *Statistical Power Analysis for the Behavioral Sciences, 2nd Edition.* Hillsdale: Lawrence Erlbaum.

Cohen, J. (1992). A power primer. *Psychological Bulletin, 112*(1), 155–159. https://doi.org/10.1037/0033-2909.112.1.155

Cormac, I., Jones, C. & Campbell, C. (2002). Cognitive behavior therapy for schizophrenia. *Cochrane Database of Systematic Reviews Database of Systematic Reviews, 2*, 1-46.

Correll, C. U., Manu, P., Olshanskiy, V., Napolitano, B., Kane, J. M., & Malhotra, A. K. (2009). Cardiometabolic risk of second-generation antipsychotic medications during first-time use in children and adolescents. *JAMA, 302*(16), 1765–1773. https://doi.org/10.1001/jama.2009.1549

Department for Education (2016). *Mental health and behaviour in schools: Departmental advice for school staff.* Retrieved from https://www.gov.uk/government/publications/mental-health-and-behaviour-in-schools--2

Department for Education and Department of Health (2015). *Special Educational Needs and Disability Code of Practice: 0 to 25 years*. Available at: https://www.gov.uk/government/publications/send-code-of-practice-0-to-25

Department of Health and Social Care, DHSC (2001). *The Mental Health Policy Implementation Guide*. London: DHSC

Dingemans, P. M., Linszen, D. H., Lenior, M. E., & Smeets, R. M. W. (1995). Component structure of the expanded Brief Psychiatric Rating Scale (BPRS-E). *Psychopharmacology, 122*, 263-267

Drake, R., Haddock, G., Tarrier, N., Bentall, R., & Lewis, S. (2007). The Psychotic Symptom Rating Scales (PSYRATS): Their usefulness and properties in first episode psychosis. *Schizophrenia Research*, *89*(1-3), 119-122.

https://doi.org/10.1016/j.schres.2006.04.024

Dunsmuir, S., & Hardy, J. (2016). *Delivering Psychological Therapies in Schools and Communities*. Leicester: BPS

Falcone, T., Mishra, L., Carlton, E., Lee, C., Butler, R. S., Janigro, D., Simon, B., & Franco, K. (2010). Suicidal behavior in adolescents with first-episode psychosis. *Clinical Schizophrenia & Related Psychoses, 4*(1), 34–40. https://doi.org/10.3371/CSRP.4.1.2

Favrod, J., Rexhaj, S., Ferrari, P., Bardy, S., Hayoz, C., Morandi, S., Bonsack, C., & Giuliani, F. (2012). French version validation of the psychotic symptom rating scales (PSYRATS) for outpatients with persistent psychotic symptoms. *BMC Psychiatry*, *12*, 161. https://doi.org/10.1186/1471-244X-12-161

Fleischhaker, C., Schulz, E., Tepper, K., Martin, M., Hennighausen, K., & Remschmidt, H. (2005). Long-term course of adolescent schizophrenia. *Schizophrenia Bulletin, 31*(3), 769–780. https://doi.org/10.1093/schbul/sbi014

Fowler, D. R., Garety, P. A., & Kuipers, E. (1995). *Cognitive behaviour therapy for psychosis: theory and practice*. Chichester: Wiley

Francey, S. M., O'Donoghue, B., Nelson, B., Graham, J. R., Baldwin, L., Yuen, H. P., Kerr, M. J., Ratheesh, A., Allott, K. A., Alvarez-Jimenez, M., Fornito, A., Harrigan, S., Thompson, A., Wood, S. J., Berk, M., & McGorry, P. D. (2020). Psychosocial Intervention With or Without Antipsychotic Medication for First-Episode Psychosis: A Randomized Noninferiority Clinical Trial. *Schizophrenia Bulletin Open, 1*(1) https://doi.org/10.1093/schizbullopen/sgaa015

Fuggle, P. & Dunsmuir, S. (2013). Working with schools and children's wider social environment. In P. Graham & S. Reynolds (Eds.), *Cognitive behaviour therapy for children and families* (3rd ed.). Cambridge: Cambridge University Press. https://doi:10.1017/CBO9781139344456.013

Fusar-Poli, P., Rocchetti, M., Sardella, A., Avila, A., Brandizzi, M., Caverzasi, E., Politi, P., Ruhrmann, S., & McGuire, P. (2015). Disorder, not just state of risk: meta-analysis of functioning and quality of life in people at high risk of psychosis. *The British journal of Psychiatry, 207*(3), 198–206. https://doi.org/10.1192/bjp.bp.114.157115

Gersten, R., Fuchs, L. S., Compton, D., Coyne, M., Greenwood, C., & Innocenti, M. S. (2005). Quality indicators for group experimental and quasi-experimental research in special education. *Exceptional Children, 71*(2), 149–164.

https://doi.org/10.1177/001440290507100202

Gough, D. (2007). Weight of evidence: A framework for the appraisal of the quality and relevance of evidence. *Research Papers in Education, 22*(2) 213-228.

Haddad, P. M., & Sharma, S. G. (2007). Adverse effects of atypical antipsychotics: differential risk and clinical implications. *CNS Drugs, 21*(11), 911–936.

Hilsenroth, M. J., Ackerman, S. J., Blagys, M. D., Baumann, B. D., Baity, M. R., Smith, S. R., Price, J. L., Smith, C. L., Heindselman, T. L., Mount, M. K., & Holdwick, D. J. Jr. (2000). Reliability and validity of DSM-IV axis V. *The American Journal of Psychiatry*, *157*(11), 1858–1863. https://doi.org/10.1176/appi.ajp.157.11.1858

Hollis C. (2000). Adult outcomes of child- and adolescent-onset schizophrenia: diagnostic stability and predictive validity. *The American Journal of Psychiatry, 157*(10), 1652–1659. https://doi.org/10.1176/appi.ajp.157.10.1652

Hui, C., Morcillo, C., Russo, D. A., Stochl, J., Shelley, G. F., Painter, M., Jones, P. B., & Perez, J. (2013). Psychiatric morbidity, functioning and quality of life in young people at clinical high risk for psychosis. *Schizophrenia Research, 148*(1-3), 175–180. https://doi.org/10.1016/j.schres.2013.05.026

Immonen, J., Jääskeläinen, E., Korpela, H., & Miettunen, J. (2017). Age at onset and the outcomes of schizophrenia: A systematic review and meta-analysis. *Early Intervention in Psychiatry, 11*(6), 453–460. https://doi.org/10.1111/eip.12412

Iyer, S. N., Mangala, R., Anitha, J., Thara, R., & Malla, A. K. (2011). An examination of patient-identified goals for treatment in a first-episode programme in Chennai, India. *Early Intervention in Psychiatry, 5*(4), 360–365. https://doi.org/10.1111/j.1751-7893.2011.00289.x

Jackson, H. J., McGorry, P. D., Killackey, E., Bendall, S., Allott, K., Dudgeon, P., Gleeson, J., Johnson, T., & Harrigan, S. (2008). Acute-phase and 1-year follow-up results of a randomized controlled trial of CBT versus Befriending for first-episode psychosis: the ACE project. *Psychological Medicine, 38*(5), 725–735. https://doi.org/10.1017/S0033291707002061

Jadad, A. R., Moore, R. A., Carroll. D., Jenkinson, C., Reynolds, D. J. M., Gavaghan, D. J., & McQuay, H. J. (1996). "Assessing the quality of reports of randomized clinical trials: Is blinding necessary?". *Controlled Clinical Trials, 17*(1), 1–12. https://doi:10.1016/0197-2456(95)00134-4

Johnson, S. B., Blum, R. W., & Giedd, J. N. (2009). Adolescent maturity and the brain: the promise and pitfalls of neuroscience research in adolescent health policy. *The Journal of Adolescent Health, 45*(3), 216–221. https://doi.org/10.1016/j.jadohealth.2009.05.016

Jones, S. R., Carley, S., & Harrison, M. (2003). An introduction to power and sample size estimation. *Emergency Medicine Journal, 20*(5), 453–458.

Lader, M. (2000). Rating scales in schizophrenia: a review of their usefulness for assessing atypical antipsychotics. *CNS Drugs, 14*(1), 23–32

Landa, Y. (1987). *Cognitive behavioral therapy for psychosis (CBTp): An introductory manual for clinicians*. United States Department of Veterans Affairs, Mental Illness, Research, Education, and Clinical Center

https://www.mirecc.va.gov/visn2/docs/CBTp_Manual_VA_Yulia_Landa_2017.pdf

Morrison, A. P. (2001). The interpretation of intrusions in psychosis: An integrative cognitive approach to hallucinations and delusions. *Behavioural and Cognitive Psychotherapy, 29*, 257-276.

Morrison, A. P. (2017). A manualised treatment protocol to guide delivery of evidence-based cognitive therapy for people with distressing psychosis: learning from clinical trials. *Psychosis, 9*(3), 271-281. DOI: 10.1080/17522439.2017.1295098

Morrison, A. P., Pyle, M., Byrne, R., Broome, M., Freeman, D., Johns, L., James, A., Husain, N., Whale, R., MacLennan, G., Norrie, J., Hudson, J., Peters, S., Davies, L., Bowe, S., Smith, J., Shiers, D., Joyce, E., Jones, W., Hollis, C., Maughan, D. (2021). Psychological intervention, antipsychotic medication or a combined treatment for adolescents with a first episode of psychosis: the MAPS feasibility three-arm RCT. *Health Technology Assessment (Winchester, England), 25*(4), 1–124. https://doi.org/10.3310/hta25040

Morrison, A. P., Pyle, M., Maughan, D., Johns, L., Freeman, D., Broome, M. R., Husain, N., Fowler, D., Hudson, J., Maclennan, G., Norrie, J., Shiers, D., Hollis, C., James, A., Morrison, A. P., Pyle, M., Maughan, D., Johns, L., Freeman, D., James, A. (2020). Antipsychotic medication versus psychological intervention versus a combination of both in adolescents with first-episode psychosis (MAPS): a multicentre, three-arm, randomised controlled pilot and feasibility study. *The Lancet Psychiatry, 7*(9), 788–800. https://doi.org/10.1016/S2215-0366(20)30248-0

Mortan Sevi, O., Tekinsav Sütcü, S., & Güneş, B. (2016). İşitsel Varsanı ve Sanrıların Değerlendirilmesi: Psikotik Belirti Değerlendirme Ölçeklerinin (PSYRATS) Türkçe Formunun Güvenilirliği ve Geçerliği [The Assessment of the Auditory Hallucinations

Symptom Rating Scales (PSYRATS)]. *Turk Psikiyatri Dergisi = Turkish Journal of Psychiatry*, *27*(2), 0.

Müller, H., Kommescher, M., Güttgemanns, J., Wessels, H., Walger, P., Lehmkuhl, G., Kuhr, K., Hamacher, S., Lehmacher, W., Müller, K., Herrlich, J., Wiedemann, G., Stösser, D., Klingberg, S., & Bechdolf, A. (2020). Cognitive behavioral therapy in adolescents with early-onset psychosis: a randomized controlled pilot study. *European Child & Adolescent Psychiatry, 29*(7), 1011–1022. https://doi.org/10.1007/s00787-019-01415-4

Murphy, A. L., Gardner, D. M., Kisely, S., Cooke, C., Kutcher, S. P., & Hughes, J. (2015). A Qualitative Study of Antipsychotic Medication Experiences of Youth. *Journal of the Canadian Academy of Child and Adolescent Psychiatry, 24*(1), 61–69.

Naeem, F., Farooq, S., & Kingdon, D. (2014). Cognitive behavioral therapy (brief vs standard duration) for schizophrenia. *Schizophrenia Bulletin, 40*(5), 958–959. https://doi.org/10.1093/schbul/sbu113

National Institute for Health and Care Excellence (2016). *Psychosis and schizophrenia in children and young people: recognition and management*. Clinical guideline CG155. https://www.nice.org.uk/guidance/cg155

Olivo, S. A., Macedo, L. G., Gadotti, I. C., Fuentes, J., Stanton, T., & Magee, D. J. (2008). Scales to assess the quality of randomized controlled trials: a systematic review. *Physical Therapy, 88*(2), 156–175. https://doi.org/10.2522/ptj.20070147

Orygen Youth Health Research Centre (2010). *Cognitive-Behavioural Case Management in Early Psychosis: A Handbook*. Melbourne, Australia: Orygen Youth Health Research Centre

Overall, J. E., & Gorham, D. R. (1962). The Brief Psychiatric Rating Scale. *Psychological Reports*, *10*(3), 799–812.

Passby, L., & Broome, M. (2017). Thought disorder. *BJPsych Advances, 23*, 321-323. DOI: 10.1192/apt.bp.116.016071.

Patterson, D. A., & Lee, M. S. (1995). Field trial of the Global Assessment of Functioning Scale--Modified. *The American Journal of Psychiatry*, *152*(9), 1386–1388. https://doi.org/10.1176/ajp.152.9.1386

Phiri, P., Rathod, S., & Carr, H. (2017). A Brief Review of Key Models in Cognitive Behaviour Therapy for Psychosis. *Acta Psychopathologica, 3*(84). Doi:10.4172/2469-6676.100156

Rabany, L., Weiser, M., Werbeloff, N., & Levkovitz, Y., (2011). Assessment of negative

Schimmelmann, B. G., Conus, P., Cotton, S., McGorry, P. D., & Lambert, M. (2007). Pre-treatment, baseline, and outcome differences between early-onset and adult-onset psychosis in an epidemiological cohort of 636 first-episode patients. *Schizophrenia Research, 95*(1-3), 1–8. https://doi.org/10.1016/j.schres.2007.06.004

Schulz, K. F., Altman, D. G., & Moher, D. (2010). CONSORT 2010 Statement: updated guidelines for reporting parallel group randomised trials. *BMJ, 340*, c332. doi: 10.1136/bmj.c332

Shaffer, D., Gould, M. S., Brasic, J., Ambrosini, P., Fisher, P., Bird, H., & Aluwahlia, S. (1983). A children's global assessment scale (CGAS). *Archives of General Psychiatry*, *40*(11), 1228-1231.

Stafford, M. R., Mayo-Wilson, E., Loucas, C. E., James, A., Hollis, C., Birchwood, M., & Kendall, T. (2015). Efficacy and safety of pharmacological and psychological interventions for the treatment of psychosis and schizophrenia in children, adolescents and young adults: a systematic review and meta-analysis. *PloS One, 10*(2), e0117166. https://doi.org/10.1371/journal.pone.0117166

Steinhausen, H. C. (1987). Global assessment of child psychopathology. *Journal of the American Academy of Child and Adolescent Psychiatry*, *26*(2), 203–206. https://doi.org/10.1097/00004583-198703000-00014

van Beek, J., Vuijk, P. J., Harte, J. M., Smit, B. L., Nijman, H., & Scherder, E. J. (2015). The factor structure of the Brief Psychiatric Rating Scale (expanded version) in a sample of forensic psychiatric patients. *International Journal of Offender Therapy and Comparative Criminology*, *59*(792 340 T0 Td [[(ne)106nal of

VandenBos, G. R. (2015). *APA dictionary of psychology*. Washington, DC: American Psychological Association

Ventura, J., Green, M. F., Shaner, A. L., & Liberman, R. P. (1993). Training and quality assurance with the Brief Psychiatric Rating Scale: "The drift busters". *International Journal of Methods in Psychiatric Research*, 34, 221–244

World Health Organization (WHO) (1992). *The ICD-10 Classification of Mental and Behavioural Disorders: Clinical Descriptions and Diagnostic Guidelines*. Geneva: WHO

Yehya, A., Ghuloum, S., Mahfoud, Z., Opler, M., Khan, A., Hammoudeh, S., Abdulhakam, A., Al-Mujalli, A., Hani, Y., Elsherbiny, R., & Al-

Sönmez, N., Romm, K. L., Østefjells, T., Grande, M., Jensen, L. H., Hummelen, B., Tesli, M., Melle, I., & Røssberg, J. I. (2020). Cognitive behavior therapy in early psychosis with a focus on depression and low self-esteem: A randomized controlled trial. Comprehensive P

Haddock, G., Tarrier, N., Morrison, A. P., Hopkins, R., Drake, R., & Lewis, S. (1999). A pilot study evaluating the effectiveness of individual inpatient cognitive-behavioural therapy in early psychosis. Social psychiatry and Psychiatric Epidemiology, 34beha4c **a**id (id)5 (e)

O'Donoghue, B., Francey, S. M., Nelson, B., Ratheesh, A., Allott, K., Graham, J., Baldwin, L., Alvarez-Jimenez, M., Thompson, A., Fornito, A., Polari, A., Berk, M., Macneil, C., Crisp, K., Pantelis, C., Yuen, H. P., Harrigan, S., & McGorry, P. (2019). Staged treatment and acceptability guidelines in early psychosis study (STAGES): A randomized placebo controlled trial of intensive psychosocial treatment plus or minus antipsychotic medication for first-episode psychosis with low-risk of self-harm or aggression. Study protocol and baseline characteristics of participants. Early Intervention in P

Weight of Evidence A: Methodological Quality Criteria

: A Jadad score of ≤2 qualifies as a low quality, a score of 3 qualifies as adequate (medium) quality, while a score of ≥4 qualifies as a high quality study.

Weight of Evidence B: Methodological Relevance Criteria

Single group design with no control group	At least one 'no intervention' comparison group (i.e. on waiting list)	A minimum of one 'active' comparison group receiving an alternative intervention or attention placebo (i.e. standard care)	This allows for an estimation of change caused by regression to the mean and the control of variables not related to the intervention (Barnett et al., 2005)
Limited or no information offered on			

	3	3	3	3	3
	3	3	3	3	3
	1	2	3	3	2
	2	3	3	3	3
	2	2	3	3	3
	2	1	2	1	2
	2.16 - Medium	2.33 - Medium	2.83 - High	2.66 - High	2.66 - High

Note ≤1.4: low
1.5 - 2.4: medium
≥2.5: high

Weight of Evidence C: Topic Relevance Criteria

The following formulae, taken from Cohen's (1988) calculations, were used to calculate effect sizes in trials where the aforementioned data was not provided.

Cohen's $d = (M_2 - M_1) / SD_{pooled}$

$SDpooled = \sqrt{[(s_1^2 + s_2^2) / 2]}$

For within group calculations where sample sizes changed over time due to participant withdrawals or drop-outs, the Psychometrica calculator was used to adjust the calculation of the pooled standard deviation with weights for the sample sizes.

d	
0.2	Small
0.5	Medium
0.8	Large