parents report school functioning problems to be their main concern (Robin, 1998). However, positive treatment in childhood has failed to produce successful long-term effects (Molina et al., 2009).

All teenagers face challenges such as managing more complex academic environments (Eccles, 2004), building increasingly complex relationships (Steinberg & Morris, 2001) and impulsivity to choose activities which incur serious negative consequences (Flory et al., 2006). Adolescents with ADHD possess difficulties that intensify these. Executive functioning difficulties i.e. organisation, time management and planning; and motivation deficits i.e. dislike for difficult tasks; are common in young people with ADHD. These may prevent teens from engaging positively with the demands of secondary school (Langberg et al., 2013). Executive functioning and motivation deficits can also exacerbate irresponsible adolescent decision-making (Casey et al., 2008). This combination makes academic functioning harder and significantly hinders the achievement of educational milestones in teenagers with ADHD (Chan et al., 2016).

2.2 Behavioural parent training (BPT) combined with organisational skills training

Historically, medication has been the main treatment in dealing with ADHD symptoms and functioning

3. CRITICAL REVIEW OF THE EVIDENCE

3.1 Literature Review

The databases PsycInfo, Medline and ERIC (EBSCO) were searched due to the review topic spanning psychology, medicine and education. They were searched on 13th January 2023. The search terms outlined in Table 1 were used. An ancestral search also generated two new articles for inclusion.

After duplicates were removed, 280 studies were screened against the inclusion and exclusion criteria outlined in Table 2, initially by title screen

Table 1

Database Search Terms

INTERVENTION		PARTICIPANT		ADHD
Behavio* parent* training OR BPT OR skill* training OR organi* skill* training OR academic training OR parent* intervention	A N D	Adolescen* OR Teen* OR "Secondary school" OR "Middle school" OR "High school"	A N D	ADHD OR "Attention Deficit Hyperactivity Disorder" OR "attention-deficit/hyperactivity disorder" OR

Table 2

Inclusion and exclusion criteria of studies for current review

Study Feature	Inclusion Criteria	Exclusion Criteria	Rationale
1) Types of publications	Peer-reviewed journal articles.	Non-peer reviewed journal articles.	Peer-reviewed journal articles have been quality assured and are therefore of a higher quality and credibility.
2) Language	Studies published in the English language.	Studies published in a language other than English.	The author of this review does not have resources for translation.
3) Country of study	Study conducted in an Organisation for Economic Cooperation and Development (OECD) country.	Studies conducted in non-OECD countries.	OECD countries are considered more comparable to a UK context because they have similar educational contexts. Non-OECD countries are not as similar as they have differing educational systems and policies.

Study Feature	Inclusion Criteria	Exclusion Criteria	Rationale
4) Research design	Empirical studies including: Randomised Control Trials (RCTs), quasi-experimental studies, single case experimental designs, cohort studies, systematic reviews	Qualitative studies, surveys	

Study Feature	Inclusion Criteria	Exclusion Criteria	Rationale
6) Diagnosis	All study participants who met the criteria for ADHD in DSM-5.	Adolescents who did not meet the criteria for ADHD.	This study is looking at the target population of those with ADHD.
7) Outcome measures	a) Studies that include at least one quantitative measure of academic functioning, such as Grade Point Average (GPA), including baseline measurements. b) A quantitative measure of a behavioural or organisational aspect of academic work including baseline measurements, such as the parent rated Homework Problems Checklist (Anesko, Schoiock, Ramirez & Levine, 1987) or teacher reported percentage of homework turned in.	Studies which do not include a measure of academic outcome, do not report the change over time, or only provide qualitative descriptions of change.	This review question is focused on the improvement of academic functioning outcomes. Exam, quiz and homework grades are a measure of this. As the intervention also aims to improve the organisational skills of adolescents, the author wanted to find studies that looked at the effect of the intervention on these skills and behaviours as well as the effect on academic grades.

Table 3

Reference list of the seven studies included in this review

Studies included in the review
Hogue, A., Fisher, J. H., Dauber, S., Bobek, M., Porter, N.,

Child & Adolescent Psychology, 49(4), 476–492.

https://doi.org/10.1080/15374416.2019.1585257

3.2 Weight Of Evidence

Gough's (2007) Weight of Evidence (WoE) Framework was used to assess the selected studies. The research was systematically evaluated, based on three dimensions: methodological quality, methodological relevance and topic relevance.

WoE A considered the methodological quality of each piece of research. For six of the studies an adapted version of Kratochwill's (2003) group intervention protocol was used. This coding protocol was considered to be the most relevant for the six RCT studies because it is very detailed in its analysis of the methodology. Five out of six of the reviewed studies were carried out by the researcher who developed the Supporting Teens' Autonomy Daily (STAND) intervention and therefore this review needed to be able to pick out detailed differences between each study. See Appendix C for details of adaptations

The ratings for WoE A, B and C were averaged to give an overall WoE D rating (Table 4). These scores were categorised into low (1.4 and below), medium (1.5 – 2.4) and high (above 2.4). WoE D calculates the overall strength of each study in answering the current review question. Appendices G and H detail the coding for each study.

Table 4

Summary of Weight of Evidence

Study	WoE A	WoE B	WoE C	WoE D
Hogue et al., 2021	0.8	1	1.5	1.1 (Low)
Raggi et al., 2009	1.7	1	1.25	1.3 (Low)
Sibley et al., 2013	1.8	2	1.5	1.8 (Medium)

3.3 Participants

Across all seven studies, 744 parents (accounted for by the number of adolescents) were involved in trials to look into the effectiveness of combined behaviour and organisational skills training. The age

was conducted in the community and trainers were community clinicians. This was reflected in a higher score in WoE C.

3.5

effects are because of the active components of the specific intervention, rather than receiving attention and expecting treatment.

3.6 Academic Functioning Measures

All seven studies used a variety of measures to assess adolescent academic functioning. Studies scored higher on WoE A for the wider range of measures and the wider source of the report e.g. school, parent and adolescent reports.

All studies provided curriculum-based measures of academic achievement. Hogue et al. (2021) used adolescent reported school grades. The researchers stated that teen self-report is reasonably valid (Crockett et al., 1987). However, other research has shown that adolescents with ADHD are notoriously bad at reporting their own functioning (Fischer et al., 1993). This was therefore considered to be a weak measure in WoE A. Sibley et al. (2014) used one-month average assignments and quiz grades whereas the other four Sibley et al. studies used teacher rated core subject GPA scores averaged across 3 months. The main issue with using average scores across multiple subjects is that the measure is unlikely to be sensitive to changes in performance over a short period of time and therefore less relevant for this review question. These measures were therefore given lower scores in WoE A. For pragmatic reasons of teacher workload, Raggi et al. (2009) only used one data point at each stage of baseline, post-treatment and follow-up. As a single-subject case design, this measure was given a lower rating in WoE A.

All seven studies measured wider academic functioning skills which are included in this review. Two studies, Hogue et al. (2021) and Raggi et al.

All training was carried out weekly for between 5 weeks (Raggi et al., 2009) and 20 weeks (Hogue et al., 2021) and sessions with specific training elements ranged from an average of 12.4 minutes per session (Hogue et al., 2021) to 90 minutes (Raggi et al., 2009, Sibley et al., 2014; 2020).

Adherence data was collected for five studies. This showed parental rated adherence to implementing strategies varied between 53% - 93%. Similarly, therapist fidelity to training varied significantly. Raggi et al. (2009) stated therapist procedural fidelity to intervention was 100%, whereas Sibley et al. (2021) ranged from 85% in skills sessions to 24% in planning sessions. And Hogue et al. (2021) had a flexible model of delivery which meant fidelity varied across all participants. The strength of training and intervention fidelity where provided, was included in WoE A ratings.

3.8 Findings and Effect Sizes

Each study's academic functioning findings were collated and effect sizes calculated to compare results. Cohen's d effect size could not be calculated for GPA scores in two studies (Hogue et al., 2021; Sibley et al., 2021) due to missing data (standard deviations or means). These studies received lower WoE A ratings. Table 5 reports the effect size descriptors to support the interpretation of these values. A summary of each of the study's academic functioning findings and effect sizes can be seen in Table 6.

Table 5

Effect size descriptors for Cohen's d, (1988)

Cohen's d	Descriptor

0.8	Large
0.5	Medium
0.2	Small

Study	Sample Size	Relevant Outcome Measure	Cohen's d	Effect size descriptor	Significance value	Overall WoE D
Sibley et al., 2013 (STAND)	1	1.GPA	0.25	Small	p<0.05*	1.8 (Medium)
		2.Academic problems (parent report)	1.30	Large	p<0.05*	
		3. Academic problems (teacher report)	0.00	None	Non-sig	
		4.Planner use	5.15	Large	p<0.05*	
Sibley et al., 2014 (STAND-G)	1 23	1.Average assignment grade				
		2.Average quiz grade				
		3.% assignments turned in aŒÌ¶â				

Study	Sample Size	Relevant Outcome Measure	Cohen's d	Effect size descriptor	Significance value	Overall WoE D
Sibley et al., 2016 (STAND)	1	1.GPA	0.31	Small	0.45	2.3 (Medium)
		2.Academic problems (parent)	1.01	Large	p<0.001*	
		3.Academic problems (teacher)	0.08	None	0.90	
		4. Homework recording	0.07	None	p 63	
Sibley et al., 2020 (STAND)	1	1.GPA	0.45	Small		
		2.Academic problems (P)	1.23	Large		
		3.Academic problems (T)	0.80			

Study	Sample Size	Relevant Outcome Measure	Cohen's d	Effect size descriptor	Significance value	Overall WoE D
Sibley et al., 2021 (STAND)	1	1.GPA	N/A[a]		N/A[a]	2.2 (Medium)
		2.Academic impairment (P)	0.57	Medium	Non-sig	
		3.Academic impairment (T)	0.24	Small	Non-sig	

Notes. The acronym CASH-AA stands for Changing Academic Support in the Home for Adolescents with ADHD

The acronym HIP stands for Homework Intervention Programme

The acronym STAND stands for Supporting Teens' Autonomy Daily

The acronym STAND-G stands for Supporting Teens' Autonomy Daily - Group

*The results reached statistical significance.

[a] Insufficient data in paper to calculate effect size or p value.

[b] No statistical analyses were conducted. Qualitative analysis provided.

[c] Effect size refers to within-group effects (i.e. pre- and post- measures of same group).

[d] Post-treatment data provided rather than follow-up

Raggi et al. (2009) collected 100% of data at baseline but only 36% of data during the follow-up phase from parents. Visual analysis of the results showed that 10 out of 11 participants showed a decline in the Homework Problems Checkm7 -2.3-6 (le)4 (m)-9 (s)-2 (Ch-(w)16 (or)72 (he r)7 (le)[0.00-2 (CC9.9u)-1 w 7.07 0 Td [(y)3 Td [(ue-1)5 (e)3 Td [(R)-3 (a)7 (g)6 (y)14 (-)1 (1)2 (0)2 (w)] TJ 7.07 0 Td [(t)

Two studies (Sibley et al., 2013; 2016) looked at specific organisational skills to support academic functioning. These both showed large effect sizes between baseline and post-treatment but this effect was not maintained 6 months later in the 2016 study. These skills were explicitly taught and so the short-term result is good. However, producing a durable outcome needs further research.

Studies analysing the factors that impact on results showed the complex nature of ADHD and working with parents and adolescents. Future studies need to look into the effects of family risk factors, as this has been highlighted as having an impact on outcomes of success (Sibley et al., 2020). Training dosages, follow-up support and therapist training level have also been highlighted (Sibley et al., 2021). ADHD presents itself in different ways and statistics suggest that between 15% and 40% (Starck et al., 2016) of parents of children with ADHD are likely to have it themselves. More research on specific elements of the programmes and how to match parent and adolescent impairment to the most appropriate intervention and support could improve results.

Overall, these study results suggest that BPT combined with organisational skills training programmes provide some benefit on the parent rating of their child's academic impairments. However, how this can be translated into an objective, durable academic impact for the adolescent, needs more research. These findings are in line with Zwi et al. (2011) who conducted a literature review on parent training interventions for children aged 5 – 18 years of age. They found mixed results between studies, with

positive results for parent training when ADHD was not comorbid with oppositional defiant disorder as well as outcomes being better for girls and for those on medication. The current studies on adolescents did not break down the results in this way, and therefore this reiterates the need for further research into who would benefit most from specific elements of this intervention.

4.2 Limitations and recommendations

A key limitation of this research was that due to the lack of studies in this area, five studies were carried out by one group of scientists, one of whom designed the treatment programme and receives royalties for publishing treatment research. This has an impact on the validity of the findings as there is a possibility of bias and conflict of interest.

Secondly, the results are difficult to generalise because the sampling of participants is very limited. All studies were carried out in the U.S.A. where the use of medication for ADHD is 10 times higher than in the U.K (Beau-Lejdstrom et al., 2016), and had unrepresentative samples. More studies need to be undertaken with larger samples, across a wider cross-section of the population and from a wider research base.

Thirdly, study conditions were highly controlled and the one study (Sibley et al., 2021) that tried to replicate community-based intervention produced poor results. More research therefore needs to be carried out in community and school settings to look at which factors best support larger numbers of parents to have a positive impact.

Finally, parents and adolescents were not blinded to conditions. This knowledge in treatment groups and control groups could have led to a change in attitude and therefore affected the self-reported levels. Future studies need to look more specifically at which elements of the programmes create positive results.

American Academy of Child and Adolescent Psychiatry, 29(4), 546–557. https://doi.org/10.1097/00004583-199007000-00007

Beau-Lejdstrom, R., Douglas, I., Evans, S. J. W., & Smeeth, L. (2016). Latest trends in ADHD drug prescribing patterns in children in the UK: Prevalence, incidence and persistence. British Medical Journal Open, 6(6), e010508. https://doi.org/10.1136/bmjopen-2015-010508

Bitsko, R. H., Claussen, A. H., Lichstein, J., Black, L. I., Jones, S. E., Danielson, M. L., Hoenig, J. M., Davis Jack, S. P., Brody, D. J., Gyawali, S., Maenner, M. J., Warner, M., Holland, K. M., Perou, R., Crosby, A. E., Blumberg, S. J., Avenevoli, S., Kaminski, J. W., Ghandour, R. M., & Meyer, L. N. (2022). Mental health surveillance among children—United States, 2013–2019. Morbidity and Mortality Weekly Report Supplements, 71(2), 1–42. https://doi.org/10.15585/mmwr.su7102a1

Casey, B. J., Jones, R. M., & Hare, T. A. (2008). The adolescent brain. Annals of the New York Academy of Sciences, 1124, 111–126. https://doi.org/10.1196/annals.1440.010

Chan, E., Fogler, J. M., & Hammerness, P. G. (2016). Treatment of Attention-Deficit/Hyperactivity Disorder in adolescents: A systematic review. Jama-Journal of the American Medical Association, 315(18), 1997–2008. https://doi.org/10.1001/jama.2016.5453

Cohen, J. (1988). Statistical power analysis for the behavioral sciences (2nd ed). L. Erlbaum Associates.

Cooper, H., Lindsay, J. J., & Nye, B. (2000). Homework in the home: How student, family, and parenting-style differences r

process. Contemporary Educational Psychology, 25(4), 464–487. https://doi.org/10.1006/ceps.1999.1036

Crockett, L. J., Schulenberg, J. E., & Petersen, A. C. (1987). Congruence between objective and self-report data in a sample of young adolescents. Journal of Adolescent Research, 2, 383–392. https://doi.org/10.1177/074355488724006

Enhancing academic and behavioral outcomes. Education & Treatment of Children, 29, 341–358.

Eccles, J. S. (2004). Schools, academic motivation, and stage-environment fit. In Handbook of adolescent psychology, 2nd ed (pp. 125–153). John Wiley & Sons, Inc.

Edwards, G., Barkley, R. A., Laneri, M., Fletcher, K., & Metevia, L. (2001). Parent–adolescent conflict in teenagers with ADHD and ODD. Journal of Abnormal Child Psychology, 29, 557–572. https://doi.org/10.1023/A:1012285326937

Evans, S. W., Schultz, B. K., DeMars, C. E., & Davis, H. (2011). Effectiveness of the Challenging Horizons After-School Program for Young Adolescents with ADHD. Behavior Therapy, 42(3), 462–474. eric. https://doi.org/10.1016/j.beth.2010.11.008

Evans, S. W., Schultz, B. K., White, L. C., Brady, C., Sibley, M. H., & Van Eck, K. (2009). A School-based organization intervention for young adolescents with Attention-Deficit/Hyperactivity Disorder. School Mental Health, 1(2), 78–88. https://doi.org/10.1007/s12310-009-9009-6

Fabiano, G. A., Pelham, W. E., Coles, E. K., Gnagy, E. M., Chronis-Tuscano, A., & O'Connor, B. C. (2009). A meta-analysis of behavioral treatments for attention-deficit/hyperactivity disorder. Clinical Psychology Review, 29(2), 129–140. https://doi.org/10.1016/j.cpr.2008.11.001

Fabiano, G. A., Pelham, W. E., Waschbusch, D. A., Gnagy, E. M., Lahey, B.

Burrows-

Journal of Clinical Child and Adolescent Psychology, 50(6), 874–887. https://doi.org/10.1080/15374416.2020.1716362

National Institute for Health and Care Excellence. (2018). Attention deficit hyperactivity disorder: Diagnosis and management [NICE Guideline No. 87]. https://www.nice.org.uk/guidance/ng87

Orvaschel, H., Puig-antich, J., Chambers, W., Tabrizi, M. A., & Johnson, R. (1982). Retrospective assessment of prepubertal major depression with the Kiddie-SADS-E.

Sonuga-Barke, E. J. S. (2003). The dual pathway model of AD/HD: An elaboration of neuro-developmental characteristics. Neuroscience and Biobehavioral Reviews, 27(7), 593–604.

6. APPENDICES

Appendix A: Table of excluded studies

Table A1

Studies that were excluded after a full text reading, with rationale

Study	Reason for exclusion (criteria number and rationale)
Evans, S. W., Schultz, B. K., Demars, C. E., & Davis, H. (2011). Effectiveness of the Challenging Horizons After-School Program for young adolescents with ADHD. Behavior therapy, 42(3), 462–474. R yrJDC 5(t)ID 5 (r)7 np68 0 T>>BDC 0 g68 0 T>>BD (h)0anJDC 5be 13.57h0 T>>2 (r)07 (y)oaT	

Molina, B. S. G., Flory, K., Bukstein, O. G., Greiner, A. R., Baker, J. L., Krug, V., & Evans, S. W. (2008). Feasibility and preliminary efficacy of an after-school program for middle schoolers with ADHD: A randomized trial in a large public middle school. Journal of Attention Disorders, 12(3), 207–217. https://doi.org/10.1177/1087054707311666	4 – This study was focused on the adolescent with a limited number of parent sessions aimed at supporting the child.
Padilla, R., & Parsons, M. H. (2019). Attention Deficit Hyperactivity Disorder Outcomes Following Remotely Administered Self-Help Training for Parents. Journal of the American Psychiatric Nurses Association, 25(5), 350–359. https://doi.org/10.1177/1078390318814616	5 – Children were aged 9 – 15 years old.
Sibley, M. H., Pelham, W. E., Evans, S. W., Gnagy, E. M., Ross, J. M., & Greiner, A. R. (2011). An evaluation of a summer treatment program for adolescents with ADHD. Cognitive and Behavioral Practice, 18(4), 530–544. https://doi.org/10.1016/j.cbpra.2010.09.002	4 – Parents were involved in a limited capacity.

Sibley, M. H., Coxe, S. J., Campez, M., Morley, C., Olson, S., Hidalgo-Gato, N., Gnagy, E., Greiner, A., Coles, E. K., Page, T., & Pelham, W. E. (2018). High versus Low Intensity Summer Treatment for ADHD Delivered at Secondary School Transitions. Journal of clinical child and adolescent psychology: the official journal for the Society of Clinical Child and Adolescent Psychology, American Psychological Association, Division 53, 47(2), 248–265. https://doi.org/10.1080/15374416.2018.1426005	4 – Parents were not involved in both groups and only in a limited capacity.
Sibley, M. H., Coxe, S. J., Zulauf-McCurdy, C., & Zhao, X. (2022). Mediators of psychosocial treatment for adolescent ADHD. Journal of consulting and clinical psychology, 90(7), 545–558. https://doi.org/10.1037/ccp0000743	This used participant data from a combination of studies already included in this review.
Steeger, C. M., Gondoli, D. M., Gibson, B. S., & Morrissey, R. A. (2016). Combined cognitive and parent training interventions for adolescents with ADHD and their mothers: A randomized controlled trial. Child neuropsychology : a journal on	8 – This study combined behavioural parent training with a computerized working memory

normal and abnormal development in childhood and adolescence, 22(4), 394–419. https://doi.org/10.1080/09297049.2014.994485

training program which is outside the remit of included interventions.

Appendix B: Mapping the field

After conducting a systematic literature search, seven studies were identified that researched the effects of behavioural parent training combined with organisational training on the academic functioning of adolescents with ADHD. The key features and differences for each of these studies are detailed in Table B1.

Table B1

Mapping the field

Authors	Location	Study Type & Control Group	Intervention Outline	Number Of Participants	Gender % And age (years)	Primary Recruit-ment criteria	Participant Characteristics	Primary Outcome Variables
Hogue, Fisher, Dauber et al. (2021)	U.S.A	Randomised control trial Comparison of two interventions	Changing Academic Support in the Home for Adolescents with ADHD (CASH-AA) vs CASH-AA + Medication Integration Protocol (MIP) Clinically flexible – average 20.5 sessions with					

38% of-17.6 (u)1023.3 ( )]T3 -4.88 -1.141 s6.6tw ih nans

Authors	Location	Study Type & Control Group	Intervention Outline	Number Of Participants	Gender % And age (years)	Primary Recruit-ment criteria	Participant Characteristics	Primary Outcome Variables
Sibley, Pelham, Derefinko, Kuriyan, Sanchez, Grazuabi (2013)	U.S.A.	Randomised control trial Treatment group assigned to STAND or a Treatment as Usual (TAU) control group.	Supporting Teens' Autonomy Daily (STAND) 8 x 60mins weekly family sessions plus optional follow up 3 x family problem solving sessions plus 1 x parent coaching session to support with teacher meeting	36 18 STAND 18 7 $ 8	72.3% male 0 H D Q D J H 12.39	Aged 11-15 Met DSM-IV-TR criteria for ADHD using		

Authors

Authors	Location	Study Type & Control Group	Specific Intervention Outline	Number Of Participants	Gender % And age (years)	Recruit-ment criteria	Participant Characteristics	Primary Outcome Variables
Sibley, Graziano, Kuriyan, Coxe, Pelham, Rodriguez, Sanchez, Derefinko, Helseth, Ward (2016)	U.S.A.	Randomised control trial assigned to STAND (treatment group) or Treatment As Usual (control group)	STAND+ motivational interviewing 10 x 50mins family sessions Optional 4 x monthly group sessions	128 adolescents and their parents 6 7 $ 1 ' 7 $ 8	64.9% male Mean age 12.75	Aged 11-15 Met DSM-IV-TR criteria for ADHD using parent interview of Computerized-Diagnostic Interview Schedule for Children (Shaffer et al., 2000) Corroborated with parent and teacher ratings of symptoms and impairment (Pelham Jr. et al., 2005) Estimated IQ of 80 or higher (WASI; Wechsler, 1999) No history of autism spectrum disorder	77.6% Hispanic, 8.8% Non-Hispanic White, 8.0% Non-Hispanic black, 5.5% Other 34.4% current medication	Parent and teacher rated Adolescent Academic Problems Checklist (AAPC). School provided Grade Point Average (GPA) % Student recorded homework Student organisation checklist (Evans et al., 2009)

Authors	Location	Study Type & Control Group	Specific Intervention Outline	Number Of Participants	Gender % And age (years)	Recruit-ment criteria	Participant Characteristics	Primary Outcome Variables
Sibley, Rodriguez, Coxe, Page and Espinal (2020)	U.S.A.	Randomised control trial Comparison of two interventions: group-based parent and adolescent skills training (STAND-G) vs individual parent and adolescent skills training (STAND)	STAND- * 8 x 90mins weekly sessions. 75mins parents only training, 15mins adolescents and parents together 6 7 $ 1 ' 10x 60mins weekly parent and adolescent sessions	123 adolescents and their parents STAND-G STAND	80.5% males 0 H D Q D J H 13.61	Aged 11-17 Met DSM-5 criteria for ADHD using parent interview of Diagnostic Interview Schedule for Children (Shaffer et al., 2000) Corroborated with teacher ratings of symptoms and impairment. No history of autism spectrum disorder or intellectual disability (IQ<70; WASI-II; Wechsler, 2011)	85.7% Hispanic, 7.9% Non-Hispanic White, 4.8%	

Authors	Location	Study Type & Control Group	Specific Intervention Outline	Number Of Participants	Gender % And age (years)	Recruit-ment criteria	Participant Characteristics	Primary Outcome Variables
Sibley, Graziano, Coxe, Bickman and Martin (2021)	U.S.A>h-6							

Appendix C: Weight of Evidence A (WoE A):

II. E – Cultural Significance

Table C2

Measurement criteria

Weighting	Criteria
Strong evidence (3)	‡ $ U H O L D E L O L W \ F R H I I L F L H Q W R I • • Collected data using multiple methods • Collected data from multiple sources
Promising evidence (2)	‡ $ U H O L D E L O L W \ F R H I I L F L H Q W • I R primary measures • Collected data using multiple methods and/or multiple sources
Weak evidence (1)	‡ $ U H O L D E L O L W \ F R H I I L F L H Q W R I • the primary outcome measures • Collected data uses single method and source
No evidence (0)	• A U H O L D E L O L W \ F R H I I L F L H Q W R I ” • Collected date from single source and/or data collected using single method.

Table C3

Comparison Group criteria

Weighting	Criteria
Strong evidence (3)	• At least one type of “active” comparison group must be used • Initial group equivalency must be established (preferably through random assignment of participants) • Evidence that change agents were counterbalanced • Less than 20% attrition.
Promising evidence (2)	• Presence of at least a “no intervention group” Evidence of at least two: • counterbalancing of change agents • group equivalence established • equivalent mortality with low attrition

Weak evidence (1)	Presence of a comparison group and at least one: • counterbalancing of change agents • group equivalence established • equivalent mortality with low attrition
No evidence (0)	• No efforts made to ensure group equivalence.

Table C4

Appropriate Statistical Analysis

Weighting	Criteria

Table C5

External Validity Criteria

Weighting	Criteria
Strong evidence (3)	• Complete and detailed description of the context within which the intervention occurs; • Provided evidence of perceived benefits from the intervention for all participant groups.
Promising evidence (2)	• Detailed description of some but not all contextual components; • Provided evidence of perceived benefits from the intervention for some participant groups.
Weak evidence (1)	• Provides overview of contextual components but lack details; • Provided evidence that participants did not perceive benefits from the intervention.

	• The analysis must provide evidence that at least 50% of the identified intervention components were necessary to produce change in the primary outcomes.
Weak evidence (1)	• Study demonstrates weak evidence for significant primary outcomes, • Study uses a design that allows for an analysis

	involving an overview of broad principles and a description of the intervention phases, or a formal or informal training session involving an overview of broad principles and a description of the intervention phases.
Weak evidence (1)	• Study must demonstrate evidence of acceptable adherence measured through at least one of the above criteria or use of a manual.
No evidence (0)	• Nothing done to ensure implementation fidelity or evidence indicates unacceptable adherence.

(cont.)

The total WoE A rating was calculated by adding together the sum of Sections A-G and dividing the scores by 7 to create an average score. The rating calculated based on these criteria is shown in Table D2. The full coding report is shown in Appendix H.

Table D2

The calculated WoE A scores and descriptors for single-subject study design, identified in the (Horner et al., 2005) protocol

Study	A Description of Participants and Setting	B Dependent Variable	C Independent Variable	D Baseline	E Experimental Control/Internal Validity	F External Validity	G Social Validity	WoE A rating
Raggi et al., 2009 (HIP)	2.7	1.8						

Appendix E: Weight of Evidence B (WoE B): Methodological Relevance

Petticrew and Roberts (2003) produced a typology of evidence to demonstrate the appropriateness of a range of methodological designs that explore the effectiveness of a particular intervention (Table E1). These criteria were used to score each study and results are shown in Table E2.

Table E1

WoE B: Coding Protocol

WoE B Rating (Qualitative Descriptor)	Design	Additional Criteria	Rationale
3 (High)	Randomised Control Trials	Random assignment to active control group. Measures taken pre/post intervention and at follow up point.	This is the hierarchy of design as set out by Petticrew and Roberts (2003). However, additional criteria have also been considered, to take into account the different strengths of the RCT studies. For each study, if any additional criteria were ticked, the lower score was given.
2 (Medium)	Cohort studies, quasi-experimental studies, single-subject case experimental designs, repeated measures design	Non-random assignment to intervention or control group. No intervention or wait-list control group. For small number designs there is data collected at least at three time points	
1 (Low)	Qualitative research, survey, non-experimental evaluation	Single group design with no allocation to groups. No comparison/control group	

		Measures taken post intervention. For small number designs there is data collected at less than three time points

Table E2

WoE B: The calculated WoE B scores and descriptors

Author	WoE B Score	Rationale
Hogue et al., 2021 (CASH-AA)	1 (Low)	No control group – within group design
Raggi et al., 2009 (HIP)	1 (Low)	School data had only one collection point
Sibley et al., 2013		

Criteria	WoE Rating and descriptor	Rationale
Intervention training provider	3.School-based clinician or trainer 2. Community clinician, trainer or therapist 1. University researcher or research assistant	The review is wanting to establish how effective the intervention would be when carried out within a wider population. A higher score was given to studies that utilised community clinicians with regular caseloads and normal supervision channels. Lower scores were given to studies that used the research director or research assistants and who had high levels of supervision as this lacks the flexibility to be widely replicated.
Relevance to the UK general population	3. Study was conducted in the UK and reported a diverse and representative sample of the population 2. Study was conducted in an OECD member country and reported a diverse and representative sample of the population or conducted in the UK but reported an	The review is looking to see how generalisable the findings are to the UK population. The countries within the Organisation for Economic Cooperation and Development (OECD, 2020) were considered to be more similar to the UK in terms of education, and

unrepresentative sample of participants.

1. Study was conducted in an OECD member country but had an unrepresentative sample of participants or was too small to be truly

Table F2

WoE C: The calculated scores and descriptors

Author	Setting replication	Intervention

Author	Setting replication	Intervention training provider	Cultural relevance to the UK	Effect Maintenance	WoE C
Sibley et al., 2021 (Community STAND)	3	2	1	0	1.5 (Medium)

Note. <1.4 is low; 1.5-2.4 is medium; >2.4 is high

Appendix G: WoE A completed coding protocols (Kratochwill, 2003)

Study 1

[Adapted from the Procedural Manual of the Task Force on Evidence-Based Interventions in School Psychology, American Psychology Association, Kratochwill, T.R. (2003)]

Coding Protocol: Comparing 2 interventions Dyadic and Group

Name of Coder: Rebecca Stokoe Date: 30/01/23

Full Study Reference in proper format: Hogue, A., Fisher, J. H., Dauber, S., Bobek, M., Porter, N., Henderson, C. E., & Evans, S. W. (2021). Randomized trial of academic training and medication decision-making for adolescents with ADHD in usual care. Journal of Clinical Child and Adolescent Psychology, 50(6), 874–887. https://doi.org/10.1080/15374416.2020.1716362

Intervention Name (description of study): Changing Academic Support in the Home for Adolescents with ADHD

☐ Type of Publication:

☐ Book/Monograph

☒ Journal Article

☐ Book Chapter

☐ Other (specify):

General Characteristics

A. General Design Characteristics

A1. Random assignment designs (if random assignment design, select one of the following)

☐ Completely randomized design

☐ Randomized block design (between participants, e.g., matched classrooms)

☐ Randomized block design (within participants)

☒ Randomized hierarchical design (nested treatments)

A2. Nonrandomized designs (if non-random assignment design, select one of the following)

☐ Nonrandomized design

☐ Nonrandomized block design (between participants)

☐ Nonrandomized block design (within participants)

☐ Prior exposure

A. Overall Rating for Measurement (select 0, 1, 2, or 3): ☐3 ☐2 ☒1 ☐0

B. Comparison Group

B1 Type of Comparison Group (Select one of the following)

☐ Typical intervention (typical intervention for that setting, without additions that make up the intervention being evaluated)

☐ Attention placebo

☐ Intervention element placebo

☒ Alternative intervention of same programme (CASH-AA + MIP)

☐ Pharmacotherapy

☐ No intervention

☐ Wait list/delayed intervention

☐ Minimal contact

☐ Unable to identify type of comparison

B2 Overall confidence of judgment on type of comparison group

☐ Very low (little basis)

☐ Low (guess)

☒ Moderate (weak inference)

☐ High (strong inference)

☐ Very high (explicitly stated)

☐ Unable to identify comparison group

B3 Counterbalancing of change agent (participants who receive intervention from a single therapist/teacher etc were counter-balanced across intervention)

☐ By change agent

☒ Statistical (analyse includes a test for intervention)

☐ Other

☐ Not reported/None

B4 Group equivalence established (select one of the following)

G1.1 Sampling procedures described in detail ☐ yes ☒ no

J5 Intensity/dosage of Intervention
☒ Unknown/insufficient information provided
☐ Information provided (if information is provided, specify both of the following:)
☐ length of intervention session:
☐ frequency of intervention session:

J6 Program Implementer (select all that apply)
☐ Research Staff
☐ School Specialty Staff
☐ Teachers
☐ Educational Assistants
☐ Parents
☐ College Students
☐ Peers
☒ Other – clinic-based therapist
☐ Unknown/insufficient information provided

J7 Training and Support Resources (select all that apply)
☒ Simple orientation given to change agents
☐ Training workshops conducted

Summary of Evidence

Indicator	Overall evidence rating 0-3	Description of evidence Strong Promising Weak No/limited evidence Or Descriptive ratings
General Characteristics		
Design		
Participants		

Appendix H: WoE A completed coding protocols (Horner et al., 2005)

Study 7

Coding Protocol : Horner, R. H., Carr, E. G., Halle, J., McGee, G., Odom, S., & Wolery, M. (2005). The use of single-subject research to identify evidence-based practice in special education, Exceptional Children, 71(2), 165-179.

Name of Coder : Rebecca Stokoe Date: 29.01.23

Full Study Reference: Raggi, V.L., Chronis-Tuscano, A., Fishbein, H. et al. Development of a Brief, Behavioral Homework Intervention for Middle School Students with Attention-

- $OO TXDOLW\ FULWHULD DUH PHW
-

- None of the quality c U L W H U L D D U H P H W

Section F: External Validity
Experimental effects are replicated across participants, settings, or materials to establish external validity.

- $ O O T X D O L W \ F U L W H U L D D U H P H W
- $ P D M R U L W \ R I T X D O L W \ F U L W H U L D D U H P H W
- A limited D P R X Q W R I T X D O L W \ F U L W H U L D L V P H W
- 1 R Q H R I W K H T X D O L W \ F U L W H U L D D U H P H W

Section G: Social Validity
The dependent variable is socially important.

- $ O O T X D O L W \ F U L W H U L D D U H P H W
- $ P D M R U L W \ R I T X D O L W \ F U L W H U L D D U H P H W
- A limited amount R I T X D O L W \ F U L W H U L D L V P H W
- 1 R Q H R I W K H T X D O L W \ F U L W H U L D D U H P H W

The magnitude of change in the dependent variable resulting from the intervention is socially important

- $ O O T X D O L W \ F U L W H U L D D U H P H W
- $ P D M R U L W \ R I T X D O L W \ F U L W H U L D D U H P H W
- $ O L P L W H G D P R X Q W R I T X D O L W \ F U L W H U L D L V P H W
- 1 R Q H R I W K H T X D O L W \ F U L W H U L D D U H P H W

Implementation of the independent variable is practical and cost effective.

- $ O O T X D O L W \ F U L W H U L D D U H P H W
- $ P D M R U L W \ R I T X D O L W \ F U L W H U L D D U H P H W
- $ O L P L W H G D P R X Q W R I T X D O L W \ F U L W H U L D L V P H W
- 1 R Q H R I W K H T X D O L W \ F U L W H U L D D U H P H W

	Overall evidence rating (0-3)	Evidence descriptors
Description of participants and setting	2.7	High
Dependent Variable	1.8	Medium
Independent Variable	2.0	Medium
Baseline	2.0	Medium
Experimental control/internal validity	1.0	Low
External validity	1.0	Low
Social validity	1.3	Low

Note. <1.4 is low; 1.5-2.4 is medium; >2.5 is high