Case Study 1: Evidence Based Practice Report

Theme: Interventions implemented by parents.

How Effective is Stepping Stones Triple P Programme for Reducing Behaviour Difficulties Among Children with Autism Spectrum Disorder?

Summary

This systematic literature review examined the effectiveness of a parenting intervention, Stepping Stones Triple P (SSTP), on reducing behavioural difficulties for children with Autism Spectrum Disorder (ASD). Through screening articles across three different databases (Web of Science, ERIC and with small to large effect sizes, indicating the effectiveness of SSTP for reducing behaviour difficulties for this population. Two of the studies received high WoE D ratings, two received medium ratings and 1 received a low rating, highlighting variation in the quality of the studies reviewed. Limitations, future directions and implications are discussed.

	Session 5: Planning ahead Session 6 -8: Implementing parenting routines Session 9: Program close

Note. This overview is based on information from Triple P (2021a); Triple P (2021b); Triple P (2021c) and Early Intervention Foundation (2023).

Psychological Basis

SSTP is a parenting intervention underpinned by social learning theory, coercion theory, behavioural family intervention and operant conditioning (Sanders, 1999).

Social Learning Theory

Social learning theory proposes that learning is a cognitive process, occurring in a social context, through observing, modelling and imitating behaviours and attitudes of others (Bandura et al., 1963, 1977). Observational learning involves observing individuals, formulating an idea of how new behaviours are performed and using this information as a guide for one's own actions. Vicarious reinforcement occurs through observation of rewards or punishments as a consequence of the behaviour (Bandura et al., 1963, 1977). SSTP is based on social learning principles, using techniques such as modelling, role play, direct feedback and home practice (Ruane & Carr, 2019).

Coercion Theory

Patterson's (1982) coercion theory, derived from behavioural research, outlines how aggressive and antisocial behaviours develop in children. Patterson (1982) proposes that ineffectual parental responses to a child's problem behaviour can unintentionally reinforce the undesired behaviour. These parent-child interactions can lead to a coercive cycle with problem behaviours increasing and escalating (Reid et al., 2002). Similar tor op r tor op 3actm. teV07 S96

A systematic literature review was conducted in December 2022, using the following databases: Web of Science, PsycINFO and ERIC (EBSCO). These databases were chosen to find articles relating to health, psychology and education. The same search terms were used across all three databases and are outlined in Table 2.

Table 2

Literature Search Terms

Search Term
"Stepping Stones Triple P" OR "SSTP" OR "Stepping Stones" OR "Triple P"
AND
"Autism Spectrum Disorder" OR ASD OR ASC OR "Autism Spectrum Condition" OR autis*

The initial search identified 114 studies. From this search, 22 duplicates were removed, leaving 92 studies. These studies were screened using the inclusion and exclusion criteria (see Table 3), resulting in 12 studies remaining. No dah 04 Tc - (e)]TJ 028 5 ET.004 T- -0.002 Tw 0.28 0 Td [(N)2 TDTc -0.0Q (

		children with ASD aged between 0-12	of children aged between 0-12	
2.	Diagnosis	Children must have a diagnosis of ASD or ASC	Children without a diagnosis of ASD or ASC	To evaluate the outcomes of this programme for children with ASD or ASC

	Criteria	Inclusion	Exclusion	Rationale
3.	Intervention	The study is based on the Stepping Stones Triple P Programme	The study does not include the Stepping Stones Triple P Programme	This review is evaluating the effectiveness of the Stepping Stones Triple P Programme
4.	Outcome	The study must include behavioural outcomes	The study does not include behavioural outcomes	This review is evaluating the effectiveness of SSTP for reducing children's behavioural difficulties
5.	Methodology	Quasi-experimental designs or randomised control trials	Studies without quasi-experimental designs or randomised control trials	Petticrew and Roberts' (2003) Hierarchy of Evidence proposes that randomised control trials, followed by quasi-experimental designs are most appropriate for evaluating effectiveness of interventions.
6.	Language	The study is published in English	The study is not published in English	Author's first language is English. Ensures the article can be fully understood and evaluated

Table 4

References of Studies Included within this Review

References
Brian, J., Tint, A., Branson., J & Pilkington., M. (2021). Effectiveness of group stepping stones positive parenting program for children with autism spectrum disorder and disruptive behaviour: Program evaluation from a large community implementation. *Journal on Developmental Disabilities*, *26*(2), 1-20.
Kasperzack, D., Schrott, B., Mingebach, T., Becker, K., Burghardt, R., & Kamp-Becker, I. (2020). Effectiveness of the Stepping Stones Triple P group parenting program in reducing comorbid behavioral problems in children with autism. *Autism the international journal of research and practice*, *24*(2), 423–436. https://doi.org/10.1177/1362361319866063

J a

Figure 1

Literature Search Process



Mapping the Field

The included studies in this review all examined the effectiveness of SSTP in reducing behavioural difficulties among children with ASD. However, they differed in the version of SSTP, research design, participants, outcome measures and implementation fidelity. In Appendix B, the key features of each study are outlined.

Weight of Evidence

Gough's (2007) Weight of Evidence (WoE) framework was used to critically appraise the five included studies, evaluating each study for quality and relevance. The WoE framework comprises of three main components; assessing methodological quality of a study (WoE A), the methodological relevance (WoE B) and the relevance of the topic to the review question (WoE C) (Gough, 2007).

For WoE A, the methodological quality of each study was assessed, using

tria(28 a(ri)3 pu)7 alEs[(K)-5(0)7 -1.728 (pr)2 (ed)10Tw 0.28 0(t)2 (ud)7 (el)6q -0.00De-002

question. The rationale and detailed criteria are shown in Appendix D for WoE B and Appendix E for WoE C. To calculate WoE D, the weightings from WoE A, B and C were averaged to give an overall weighting, outlined in Table 5.

Table 5

Weight of Evidence Ratings

Study	WoE A: Methodological quality	WoE B: Methodological relevance	WoE C: Topic relevance	WoE D: Overall weight of evidence
Brian et al. (2021)	1	1	2.25	1.42 (low)

children are suggested to prevent behavioural problems and improve wellbeing (Sanders et al., 2014).

All children had a confirmed diagnosis of ASD from a paediatrician or other medical professional. Three of the studies (Kasperzack et al., 2020;

Intervention

Four of the studies (Brian et al., 2021; Kasperzack et al., 2020; Tellegen et al., 2014; Zand et al., 2018) examined the effectiveness of SSTP as the primary intervention with two of the studies (Tellegen et al., 2014; Zand et al., 2018) using the Primary Care SSTP version and two of the studies (Brian et al., 2021; Kasperzack et al., 2020) utilising the group format of SSTP.
T

Kasperzack et al. (2020) reported a strong internal consistency score for the DBC (α = 0.94), SRS (α = 0.91), ECBI (Intensity scale: α = 0.94; Problem scale: α = 0.95) and total SDQ problem score (α = 0.82). A test-retest reliability score for DBC subscales was reported to be sufficient (ranging from .76 to .96). Similarly, Tellegen et al. (2014) reported internal consistency scores for the ECBI (Intensity scale = .91; Problem scale =.89). Through using multiple, reliable assessment tools and sources for data collection, this increases reliability and validity and consequently generalisability of results DB of s

outcome measures at two time points (pre and post intervention) and therefore received a lower rating for WoE A and C.

Implementation Fidelity

Implementation fidelity was assessed across all five studies and scored, forming part of the WoE A weighting. Three of the studies (Tellegen et al., 2014; Whittingham et al., 2009; Zand et al., 2018) received a ‘strong evidence’ WoE A ‘implementation fidelity’ rating because the practitioners had received formal SSTP training and received supervision throughout the programme to ensure the programme was implemented correctly. Kasperzack et al. (2020) and Brian et al. (2021) received a ‘weak evidence’ rating of 1 due to having received training for SSTP but not receiving supervision. Implementation fidelity is important for ensuring the programme is implemented correctly to determine if the intervention is responsible for the outcomes reported (Kratochwill, 2003).

Findings

All studies reported effect sizes, with two reporting Cohen’s *d* (Brian et al., 2021; Tellegen et al., 2014) and three Tf 0 Tc 0 Tw 12.62 0 Td 5

Table 6
Summary of Effect Sizes

Study	Outcome Measure	Effect Size (Cohen's *d*)	Descriptor	Significance (p value)	Main Findings	WoE D
Brian et al. (2021)	SDQ Total Difficulties	d=0.43				

Discrepancy was found between studies regarding the statistical analysis. Two of the studies (Brian et al., 2021; Whittingham et al., 2009) received a ‘high’ weighting on the WoE A ‘statistical analysis’ criteria, having used appropriate statistical analysis, controlling for familywise error rate through applying Bonferroni correction and having a sufficiently large sample size. Tellegen et al. (2014) used appropriate unit of analysis for all three measures and had a sufficiently large sample size, indicating promising evidence. However, familywise error rate was not shown to be controlled for in the study. Kazperzack et al. (2020) used appropriate unit of analysis for all

e i ng pros n4bJ EMC ET /Artifact B EMC /MC

2014). This reduces the ecological validity of the findings and makes it difficult to generalise these findings to other settings, such as school. Future research should utilise more information sources, such as teachers, and more assessment tools, to examine the impact of SSTP on behavioural difficulties in other settings.

Furthermore, two of the studies did not include control groups, making it difficult to conclude whether the impact was due to the intervention or confounding variables that the study did not control for which could overestimate the effectiveness of the intervention (Marsden & Carole, 2012). Therefore, more randomised control trials would strengthen the argument for the effectiveness of SSTP for reducing behaviour difficulties among children with ASD.

Three of the studies (Kasperzack et al., 2020;

References

Hatfield, D. R., & Ogles, B. M. (2004). The use of outcome measures by psychologists in clinical practice. *Professional Psychology: Research and Practice*, *35*(5), 485–491. https://doi.org/10.1037/0735-7028.35.5.485

Kasperzack, D., Schrott, B., Mingebach, T., Becker, K., Burghardt, R., & Kamp-Becker, I. (2020). Effectiveness of the Stepping Stones Triple P group parenting program in reducing comorbid behavioral problems in children with autism. *Autism : the international journal of research and practice*, *24*(2), 423–436. https://doi.org/10.1177/1362361319866063

Kratochwill, T. R. (2003). Task Force on Evidence Based Interventions in School Psychology. American Psychological Association.

Lenhard, W. & Lenhard, A. (2016) *Computation of effect size.* Retrieved from: h

setting. *European Journal of Special Needs Education*, *35*(1), 40–54. https://doi.org/10.1080/08856257.2019.1607659

Patterson, G. R. (1982). *Coercive family process*. Eugene, OR: Castalia.

Petticrew, M., & Roberts, H. (2003). Evidence, hierarchies, and typologies: Horses for courses. *Journal of Epidemiology and Community Health*, *57*, 527–529. doi:10.1136/jech.57.7.527

Reid, J. B., Patterson, G. R., & Snyder, J. J. (2002). *Antisocial behavior in children and adolescents: A developmental analysis and model for intervention*. American Psychological Association. https:// doi.org/10.1037/10468-000

Ruane, A. and Carr, A. (2019), Systematic Review and Meta-analysis of Stepping Stones Triple P for Parents of Children with Disabilities. Family Process., 58: 232-246. https://doi.org/10.1111/famp.12352

Sanders M. R. (1999). Triple P-Positive Parenting Program: towards an empirically validated multilevel parenting and family support strategy for the prevention of behavior and emotional problems in children. *Clinical child and family psychology review*, *2*(2), 71–90. https://doi.org/10.1023/a:1021843613840

Sanders, M. R., Mazzucchelli, T. G., & Studman, L. J. (2004). Stepping Stones

Steenfeldt-Kristensen, C., Jones, C. A., & Richards, C. (2020). The prevalence of self-injurious behaviour in autism: A meta-analytic study. *Journal of Autism and Developmental Disorders*, *50*(11), 3857–3873. https://doi.org/10.1007/s10803-020-04443-1

Tellegen, C. L., & Sanders, M. R. (2014). A randomized controlled trial evaluating a brief parenting program with children with autism spectrum disorders. *Journal of Consulting and Clinical Psychology, 82*(6), 1193–1200. https://doi.org/10.1037/a0037246

Triple P (2021a). *Group Stepping Stones Triple P*. Triple P International Pty Ltd.

Triple P. (2021b). *Primary Care Stepping Stones Triple P*. Triple P International Pty Ltd.

Triple P (2021c). *Standard Stepping Stones Triple P*. Triple P International Pty Ltd.

Weiss, J. A., Cappadocia, M. C., MacMullin, J. A., Viecili, M., & Lunsky, Y. (2012). The impact of child problem behaviors of children with ASD on parent mental health: The mediating role of acceptance and empowerment. *Autism,* 16(3), 261-274. https://doi.org/10.1177%2F1362361311422708

Whittingham, K., Sofronoff, K., Sheffield, J., & Sanders, M. R. (2009). Stepping Stones Triple P: an RCT of a parenting program with parents of a child diagnosed with an autism spectrum disorder. *Journal of abnormal child psychology*, *37*(4), 469–480. https://doi.org/10.1007/s10802-008-9285-x

Yates, K., & Le Couteur, A. (2016). Diagnosing autism/autism spectrum disorders. *Paediatrics and Child Health*, *26*(12), 513–518. https://doi.org/10.1016/j.paed.2016.08.004

Zand, D.H., Bultas, M.W., McMillin, S.E., Halloran, D., White, T., McNamara, D. and Pierce, K.J. (2018), A Pilot of a Brief Positive Parenting Program on Children Newly Diagnosed with Autism Spectrum Disorder. *Family Process*., 57: 901-914. https://doi-org.libproxy.ucl.ac.uk/10.1111/famp.12334

Appendices

Appendix A: Excluded Studies at Full Paper Review

Table 7

Excluded Studies

Reference	Reason for Exclusion	Inclusion/Exclusio

Reference	Reason for Exclusion	Inclusion/Exclusion Criteria Number
Schrott, B., Kasperzack, D., Weber, L., Becker, K., Burghardt, R., & Kamp-Becker, I. (2018). Effectiveness of the stepping stones triple P group parenting program as an additional intervention in the treatment of autism spectrum disorders: Effects on parenting variables. *Journal of Autism and Developmental Disorders*, *49*(3), 913–923. https://doi.org/10.1007/s10803-018-3764-x	4	Did not measure child behavioural outcomes
Sofronoff, K., Jahnel, D., & Sanders, M. (2011). Stepping stones triple P seminars for parents of a child with a disability: A randomized controlled trial. *Research in Developmental Disabilities*, *32*(6), 2253–2262. https://doi.org/10.1016/j.ridd.2011.07.046	2	Included participants without ASD
Whittngham, K., Sofronoff, K., & Sheffield, J. (2006). Stepping stones triple P: A pilot study to evaluate acceptability of the program by parents of a child diagnosed with an autism spectrum disorder. *Research in Developmental Disabilities*, *27*(4), 364–380. https://doi.org/10.1016/j.ridd.2005.05.003		

Appendi x B: Mapping the Field

Table 8

Author	Participants	Demographics	Study Design	Intervention	Outcome Measures	Outcome	Follow up
Kasperzack et al. (2020)	Parents of 24 children with ASD aged between 3.6 and 12 years	Gender of children: 21 male, 3 female Relationship to child: 20 mothers, 3 fathers No ethnicity data	Single group repeated measures design	Group SSTP 6 weeks	Developmental Behavior Checklist (DBC) (filled in by parents and teachers) Social Responsiveness Scale (SRS) (filled in by parents and teachers) Eyeberg Child Behavior Inventory (ECBI) (filled in by parent only) SDQ (filled in by teachers and parents)	Comorbid behavioural problems, measured with the parent and teacher ECBI, SDQ and SRS showed a significant decline at post-treatment DBC indicated a significant reduction in behaviour problems, measured by parent ratings but not by teacher ratings	Post-hoc analysis revealed significant effects were maintained at follow up

Author	Participants	Demographics	Study Design	Intervention	Outcome Measures	Outcome	Follow up

Author	Participants	Demographics	Study Design	Intervention	Outcome Measures	Outcome	Follow up
Zand et al. (2018)	21 parents/carers with children with newly diagnosed ASD, ages 2-12 Intervention group=12 Control group=9	Gender of child: 18 male 3 female Race of children: Caucasian= 14 African-American=5 Bi-racial=2 Gender of parent/carer: 21 female	Pretest/posttest two group design with random assignment to the PC SSTP intervention versus the Wait List Control	Primary Care SSTP (4 session manualised positive parenting programme)	ECBI	Clinically and statistically significant reductions in; intensity of child disruptive behaviours but not the number of behaviours	No follow-up

	Removed Item	Rationale
III.	Key Features for Coding Studies and Rating Level of Evidence/Support. A	

	Removed Item	Rationale
VI.	E. Cultural Significance	Not relevant to the scope of this review.
VII.	F. Educational/Clinical Significance	This is discussed elsewhere in this review.
VIII.	G. External Validity Indicators	This is discussed elsewhere in this review.
IX.	H. Durability of Intervention and Outcomes. Removed: H2 Durability/Generalization over time H3 Durability/Generalization across settings H4 Durability/Generalization across persons	This is discussed elsewhere in this review.
X.	I Identifiable Intervention Components (all sections)	Not relevant to this review.
XI.	J Implementation Fidelity: J4 Implementation Context (removed)	Not relevant to this review.
XII.	K. Replication (all sections)	Not relevant to this review.
XIII.	L. Site of Implementation	Not relevant to this review.

Table 10

Weight of Evidence (WoE) A ratings

Study	Measures	Comparison Group	Analysis	Follow up	Implementation Fidelity	Overall WoE A	Descriptor
Brian et al. (2021)	1	0	3	0	1	1	Low
Kasperzack et al., (2020)	3	0	2	3	1	1.8	Medium
Tellegen et al., (2014)	3	3	2	3	3	2.8	High
Whitting ham et al., (200367.92 602.0(00o0(.32 T8.8 519.72 Tm [(e)-2 (t a)-2 (l)4 (.,)72 Tm [(e).68 55.2 r5194							

Appendix D WoE B Methodological Relevance

The methodological relevance of each study was assessed and weighted for WoE B, using Petticrew and Roberts' (2003) Hierarchy of Evidence. Petticrew and Roberts (2003) propose that Randomised Control Trial designs are the most appropriate design for answering 'effectiveness' questions, followed by quasi-experimental designs. Studies with a control group are argued to be of higher quality compared to studies without a control group. The criteria for WoE B ratings can be found in Table 11. A rating for the quality of evidence was given for each study, with higher quality of evidence receiving a higher rating (see Table 12).

Table 11

Criteria for WoE B Ratings

Criteria	WoE Rating	Descriptor	Rationale
A: Study Type	3 (high)	Randomised Control Trial	It is important to consider the study design when assessing the quality of each study
	2 (medium)	Quasi-experimental designs with a control group	
	1 (low)	Quasi-experimental designs without a control group, cohort studies	

Table 12

WoE B Ratings for Studies in this Review

Study	WoE B	Descriptor
Brian et al. (2021)	1	Low
Kasperzack et al., (2020)	1	Low
Tellegen et al., (2014)	3	High
Whittingham et al., (2009)	3	High
Zand et al., (2018)	3	High

Note. <1.7= low, 1.7-2.4=medium, >2.4= high

Appendix E: WoE C Topic Relevance

WoE C assessed how appropriate the design of the study was to the review question. The criteria in table 13 was developed by the reviewer, considering the intervention, participant diagnosis, participant age and outcome measures. The rationale for each criteria is described in Table 13 and WoE C ratings for each study can be found in Table 14.

Table 13

Criteria for WoE C Ratings

Criteria	WoE Rating

	Criteria	WoE Rating	Descriptor	Rationale
C.	Participant Age	3		

pant

Appendix F: Coding Protocols

[Adapted from the Procedural Manual of the Task Force on Evidence-

A2. Nonrandomized designs (if non-random assignment design, select one of the following)

☐ Nonrandomized design

C. Stage of Progra m

☐ Model/demonstration programs

☐ Early stage programs

☒ Established/institutionalized programs

☐ Unknown

D. Concurrent or Historical Intervention Exposure

☒ Current exposure

☐ Prior exposure

☐ Unknown

2. Key Features for Coding Studies and Rating Level of Evidence/Support

(Rating Scale : 3= Strong Evidence, 2=Promising Evidence, 1=Weak Evidence, 0=No Evidence)

A. Measurement (Estimating the quality of the measures used to establish effects)

A1 The use of the outcome measures produce reliable scores for the majority of the primary outcomes

☒Yes

☐No

☐Unknown/unable to code

A2 Multi-method (at least two assessment methods used)

☒ Yes

☐ No

☐ N/A

☐ Unknown/unable to code

A3 Multi-source (at least two sources used self-reports, teachers etc.)

☒ Yes – teacher and parent

☐ No

☐ N/A

☐ Unknown/unable to code

A4 Validity of measures reported (well-known or standardized or norm-referenced are considered good, consider any cultural considerations)

☒ Yes validated with specific target group

☐ In part, validated for general population only

☐ No

☐ Unknown/unable to code

Overall Rating for measurement=3

3= Strong Evidence 2=Promising Evidence 1=Weak Evidence
0=No Evidence

B. Comparison Group

B1 Type of Comparison Group (Select one of the following)

☐ Typical intervention (typical intervention for that setting, without additions that make up the intervention being evaluated)

☐ Attention placebo

☐ Intervention element placebo

☐ Alternative intervention

☐ Pharmacotherapy

☐ No intervention

☐ Wait list/delayed intervention

☐ Minimal contact

☒ Unable to identify type of comparison

B2 Overall confidence of judgment on type of comparison group

☐ Very low (little basis)

☐ Low (guess)

H1.1 Timing of follow up assessment: ☒yes ☐ no

Specify – 6-month follow up

H1.2. Number of participants included in the follow up assessment: ☒yes ☐ no

Specify 22 families

H1.3, Consistency of assessment method used: ☒yes ☐ no

Summary of Evidence

Indicator	Overall evidence rating 0-3 NNR= No numerical rating	Description of evidence Strong Promising Weak No/limited evidence Or Descriptive ratings
General Characteristics		
Design	NNR	
Type of programme	NNR	
Stage of programme	NNR	

Measurement	3	Strong
Comparison group	0	No evidence