Case Study 1: An Evidence-Based Practice Review Report

Theme: School (Setting) based interventions for children with Special Educational Needs (SEN)

How Effective is Computer-Based Working Memory Training (CWMT) at Reducing Behavioural Symptoms of Attention Deficit Hyperactivity Disorder (ADHD) in Children?

1.0. Summary

Computer-based working memory training (CWMT) has evolved since Klingberg et al.'s (2002) landmark feasibility study into its use. CWMT aims to reduce the symptoms of Attention Deficit Hyperactivity Disorder (ADHD) by increasing children's working memory (WM) capacity.

percent of adults in the United Kingdom (UK), and is four times more prevalent in boys than girls (CAMHS Advisory Group, 2018).

Children with acute symptoms of ADHD are often prescribed mild psychoactive drugs which are effective for most in the short term (Faraone & Buitelaar, 2010). However, their long-term effects are unknown, there is an increased likelihood that these children will go on to abuse drugs and alcohol (Groenman et. al., 2013), and knowing the right time to stop medicating is problematic (van de Loo-Neus et al., 2011). Furthermore, the life-time prognosis for children with ADHD is concerning, as they exhibit higher incidences of "psychiatric outcomes including markedly elevated rates of antisocial, addictive, mood and anxiety disorders" (Biederman et al., 2006, p167). Consequently, there have been attempts to find non-pharmaceutical

include inhibitory EF, timing and working memory networks (Sonuga-Barke et al., 2014), although it should be noted that EF deficits are not consistently represented across ADHD populations (Barkley & Murphy, 2010). Motivational networks affected include intrinsic motivation and reward-punishment networks. Energetic processes, such as effort and arousal have also been implicated (Sonuga-Barke et al., 2014).

2.2. Working Memory

2.2.1. Psychological Theory of Working Memory

According to the Baddeley and Hitch model (1974; as cited in Lieberman, 2012), WM has three components; the phonological loop (e.g. for processing audio for about two seconds of spoken material), the visuo-spatial sketchpad (e.g. for processing images) and the central executive, which controls what information is processed, and combined, with information retrieved from long-term memory (LTM). The central executive coordinates information flow from the two lower components (Baddeley et al., 1991).

Thus, WM can be defined as "a store where we can combine information from the phonological loop, the visuo-spatial sketchpad and the long-term memory" (Lieberman, 2012, p332), under the control of the central executive. Put more simply, WM is the ability to hold and process information in the short term before integrating it with and committing it to LTM.

2.2.2. Working Memory and ADHD

The information stored in the WM is transient, and so it is sometimes referred to as short-term memory (STM). Vocabulary development (Gathercole et al., 1999) and early stage word learning (Jackson et al., 2016) were shown to be

strongly associated with phonological STM capacity. This has implications for children with ADHD and may explain why they often experience reading and academic difficulties.

Our visual-spatial memory of pictures is much more developed and effective than the phonological loop responsible for the memory of words. This is known as the 'picture superiority effect' (Defeyter et al., 2009) and may explain why children with ADHD have difficulties with mathematics, planning, deferred gratification and other abstract activities which involve visualisation.

Baddeley et al. (1991) provided evidence for the central executive controlling the phonological loop and the visuo-spatial sketchpad, with the central executive being required to coordinate activity between the two components of WM. Consistent with a central executive or EF deficit is that children with ADHD are often impulsive, inattentive and restless. Thus, deficits in all three components of WM link individually and collectively to symptoms of ADHD.

2.3. Computer-Based Working Memory Training (CWMT) Intervention

2.3.1. Psychological Theory of the Intervention

CWMT aims to take advantage of brain plasticity and increase WM capacity in children with ADHD using adaptive computer programs to either increase the efficiency of pre-cortical and parietal domains, or increase their neural volume. The theory is akin to working out in a gym to increase muscle strength: neural networks are more likely to develop using adaptive programs which respond to user input and maintain neural network operation at full capacity, rendering the need for them to either become more efficient or increase their volume to meet the demand.

According to the Baddeley and Hitch model (1974; as cited in Lieberman, 2012), with increased WM capacity, children with ADHD should have increased EF, phonological and visuospatial capabilities. This should result in better social and emotion regulation and thus relationships with others, increased focus in the classroom and overall improvements in behaviour.

2.3.2. The Basis for Change

Cogmed is a CWMT program consisting of brain-training computer games designed to increase children's WM capacity. The embedded activities include inhibition, visuospatial and forward and backward recall tasks (Klingberg et al., 2005). It is a five-week program involving five days training per week for approximately 40 minutes per day where the trainee completes 90 trials which focus on various components of working memory (Roche & Johnson, 2014). Children are supervised during training and so an adult, usually a parent, is also trained in software use. Internet access is required as training scores are uploaded to a server where the trainee's progress is monitored. Ideally, monitoring should be conducted by a Cogmed certified coach who not only trains the adult and the child, but who communicates with the adult at least weekly to ensure compliance and engagement with the programme, and addresses any technical issues or queries that may arise (Roche & Johnson, 2014).

The Cogmed programme is adaptive, and so the level of challenge adjusts in response to trainee input so that a significant level of challenge is maintained. This is to sustain motivation and optimise WM development. There are also reward games built into the programme that can be earned to

motivate trainees to complete sessions. Cogmed is considered suitable for children with ADHD because the adaptive nature, game element and embedded rewards are designed to engage children and maintain their focus, which is usually a challenge for children with ADHD, whilst increasing their WM capacity.

A 'non-adaptive' version of the Cogmed software also exists which can be used in research as an intervention element placebo. The non-adaptive version contains similar games, but the challenge is maintained at a low level, not increasing in response to the user's improved competence.

By changing brain physiology, the adaptive version of Cogmed aims to develop positive near- and far-transfer effects. The former describes transfer of trained WM capabilities to similar untrained tasks, such as visuospatial tasks, and the latter describes transfer of trained WM capabilities to dissimilar, untrained tasks, such as the behavioural symptoms of ADHD. These would manifest as increased focus in lessons, for example, being able to sit still for longer periods of time, reduced restlessness and impulsivity, and improved emotion regulation resulting in moim (as)4 (k)4 (s)4 (,)2 (s7) in l,s4 (w)p10 (m)-3

EBSCO), ProQuest, PubMed, PsychInfo and Web of Science Core Collection (WoS).

Six articles were retained for inclusion in the review from 227 results. The research journey was documented (Figure 1), and the inclusion and exclusion criteria summarised (Table 2). Proquest proved the least

Table 2

To show Inclusion and Exclusion criteria for review articles

Table 3

The Six Articles Included in the Review After Full-Text Screening

Study Code	Reference
1	Bigorra, A., Garolera, M., Guijarro, S., & Hervas, A. (2016). Long-term far-transfer effects of working memory training in children with ADHD: A randomized controlled trial. *European Child and Adolescent Psychiatry, 25,* 853-867. https://doi.org/10.1007/s00787-015-0804-3
2	Chacko, A., Bedard, A. C., Marks, D. J., Feirsen, N., Uderman, J. Z., Chimiklis, A., Rajwan, E., Cornwell, M., Anderson, L., Zwilling, A., & Ramon, M. (2014). A randomized clinical trial of Cogmed Working Memory Training in school-age children with ADHD: A replication in a diverse sample using a control condition. *Journal of Child Psychology and Psychiatry, 55*(3), 247-255. https://doi.org/10.1111/jcpp.12146
3	Green, C. T., Long, D. L., Green, D., Losif, A-M, Dixon, J. F., Miller, M. R., Fassbender, C. & Schweitzer, J. B. (2012). Will working memory training generalize to improve off-task behaviour in children with Attention-Deficit/Hyperactivity Disorder? *Neurotherapeutics, 9,* 639-648. https://doi.org/10.1007/s13311-012-0124-y
4	Hovic, K. T., Saunes, B-K., Aarlien, A. K. & Egeland, J. (2013). RCT of working memory training in ADHD: Long-term near-transfer effects. *PLOS One, 8*(12), 1-9. https://doi.org/10.1007/s13311-012-0124-y
5	Klingberg, T., Fernell, E., Olesen, P. J., Johnson, M, Gustafsson, P., Dahlstrom, K., Gillberg, C. G., Forssberg

The medication status of participants was also heterogenous with three studies excluding medicated children (Bigorra et al., 2016; Klingberg et al., 2005; van Dongen-Boomsma et al., 2014), two studies including medicated children (Green et al., 2012; Hovik et al., 2013), and one study not stating the medication status of the children (Chako et al., 2014). Klingberg et al. (2005) purposely did not recruit medicated children because they predicted that the children's ADHD symptoms would be more acute than those of unmedicated children, and their progress less evident as a result of intervention.

3.2.3. The Cogmed Intervention

All studies used the Cogmed CWMT program, with one study using the junior version (van Dongen-Boomsma et al., 2014) and one employing Intent to Treat (ITT) instead of non-adaptive Cogmed (Hovik et al., 2013).

Table 4

Mapping the field to present an overview of the studies included in the review

Study Number/ Reference	Country of Origin	Age Group (Years)	Number of Participants	Gender %		ADHD Diagnosis	Medication Accepted?	Computer Programme Used	Control Group Activity
				Boys	Girls				
1. Bigorra et al. (2016)	Spain	7-12	66	45	55	*DCM-IV combined Type ADHD	No medication	RoboMemo Cogmed	Non-adaptive RoboMemo

3.3. Weight of Evidence (WoE)

medication for ADHD (*n*=8) than the comparison group (*n*=2)" (Green et al., 2012, p643), rendering the groups unbalanced.

Only one study (Bigorra et al., 2016) included a power calculation, and three studies offered insufficient or no rationale for the sample size used (Green et al., 2012; Hovik et al., 2013; Klingberg et al., 2005). Furthermore, Chacko et al. (2014, p251) input missing data "with the assumption that the missing data are at least missing at random", which may have altered the outcomes, considering the impact of attrition of nine out of 85 children. This comprised 11 percent of the sample, and suggested that the study was under-powered.

3.3.2.2. Participant Recruitment

parents and children in the treatment (adaptive) group was proposed to inflate parent ratings on questionnaires (Chacko et al., 2014).

Consequently, Chacko et al. (2014) focused on ensuring that the conditions of the non-adaptive group, such as equal time spent on training tasks, mirrored those of the adaptive group, and participants were provided with extra coaching support and intervention to modify training tasks according to their needs. However, in terms of Methodological Relevance (WoE B; Appendix E, Table E), Chacko et al. (2014) and Green et al. (2012) were both judged Promising, whilst Klingberg et al. (2005) was judged Weak, indicating that the anticipated improvements in Chacko et al.'s (2014) methodology did not manifest in terms of their relevance to the review question.

3.3.3. Study Relevance to the Review Question (WoE C)

The effectiveness of the studies to ascertain if CWMT improves behavioural outcomes for children with ADHD was varied (Appendix F, Table G). Bigorra et al. (2016) was the only Strong study in this area, and notably, the only study demonstrating an overall significant impact of CWMT on behavioural ADHD outcomes (Table 5). Two studies were judged Promising (Chacko et al., 2014; van Dongen-Boomsma et al., 2014) and two were judged as Weak; Green et al. (2012) and Klingberg et al. (2005). Hovic et al. (2013) measured WM but not far transfer effects such as behavioural outcomes, and so was judged as providing No/Limited evidence of study relevance to the review question.

3.3.3.3. Confirmation of ADHD Status

Overall, confirmation of ADHD status was Promising, with some variation between studies (Table 4). The Strong studies recruited participants from establishments where they were already diagnosed with ADHD (Bigorra et al., 2016; Hovic et al., 2013), so more was known about the children's condition. In the only Weak study, participants were recruited from the community and non-clinical "consensus diagnosis based on parent and teacher ratings" used to assign ADHD status (Chacko et al., 2014, p249).

3.3.4. Overall Review Judgment (WoE D)

The review was judged Promising overall (Appendix G), however, no areas were judged Strong. Methodological Quality (WoE A) was Promising, highlighting that reviewing RCTs does not guarantee encountering high quality studies, although the structure embedded in RCTs and the Cogmed intervention may have inflated th). 2 (e Cyi (Cu)16(as2,3 T4[(t)2e(4d(t)20(es)4-b6(1)(4a(n)(4e(

(BRIEF; Gioia et al., 1996), with the caveat that only the significant results were reported and so an overall effect size could not be calculated. The two Weak studies according to WoE D generated mixed results with large (off-task behaviour) and small (fidgets) effect sizes rendering the overall result inconclusive (Green et al., 2012), and medium (hyperactivity/impulsivity) and small (inattention) effect sizes which were considered with caution due to the poor methodology employed (Klingberg et al., 2005). Moreover, the effect sizes calculated from the raw data published by Klingberg et al. (2005), using the Campbell Calculator (Wilson, 2022), were lower than those published in the article, which cited "corrected values" without a clear explanation of their rationale for correction (Klingberg et al., 2005, p181). Furthermore, it was noted that Klingberg et al. (2005), used one-tailed tests during statistical analyses, which reduced the statistical power required for significant outcomes, and justified this by stating that the study was based on previous research findings (Klingberg et al., 2002). Thus, the two Weak studies, plus a third study which did not measure behavioural outcomes (Hovic et al., 2013), contributed little to the overall review conclusions.

The remaining two studies were both judged Promising (WoE D) and both reported little or no effect of CWMT on behavioural symptoms of ADHD (Chacko et al., 2014; van Dongen-Boomsma et al., 2014). Interestingly, the latter used the BRIEF questionnaire (Gioia et al., 1996), as did Bigorra et al. (2016), but reported an effect size for the whole questionnaire rather than just significant elements. This may explain why one result supported CWMT (Bigorra et al., 2016) and the other did not (van Dongen-Boomsma et al., 2014) despite the same questionnaire being used. Alternatively, the contrast

may be due to the participants' age (5-7 years old verses 7-12 years old), sex (more boys than girls verses more girls than boys), or the computer program used (Cogmed Junior verses RoboMemo Cogmed), and highlighted

chal2p (s)4 (es)4 (R)6 (oboM)7 (em)hTd6 (am)]Tn2eBd[(c)4

Table 5

Summary of the review findings

Study	Sample Size	Primary Outcome Measure(s)	Post-Intervention and Follow-Up Effect Sizes	Effect Size Descriptor	Does CWMT Improve behavioural ADHD Symptom's in Children?	WoE D
1. Bigorra et al. (2016)	66 (Active n=36) (Control n=30)	[2]BRIEF-Parent T2 = 6-month follow up	Working Memory Subscale T2-T1 *d*= -0.86 T2-T0 *d*= -0.61 Plan/Organise T2-T1 *d*= -0.71 Metacognition Index T2-T1 *d*= -0.78	 Large Medium Medium Medium	Small to Large Effect though most effects were Medium	2.62 Strong
		[2]BRIEF-Teacher T2 = 6-month follow up	Initiate Subscale T1-T0 *d*= -0.55 T2-T0 *d*= -0.57 Working Memory T1-T0 *d*= -0.36 T2-T0 *d*= -0.84 Metacognitive Index T1-T0 *d*= -0.37 T2-T0 *d*= -0.81 Monitor Subscale T2-T1 *d*= -0.72 T2-T0 *d*= -0.79	 Medium Medium Small Large Small Large Medium Medium		

Study	Sample Size	Primary Outcome Measure(s)	Post-Intervention and Follow-Up Effect Sizes	Effect Size Descriptor	Does CWMT Improve behavioural ADHD Symptom's in Children?	WoE D
			Shift Subscale T2-T1 *d*= -0.39	Small		
2. Chacko et al. (2014)	85 (Active n=44) (Control n=41)	Parent				

Study	Sample Size

4.0. Conclusion and Recommendations

The findings of the strongest study in the sample indicated that CWMT significantly improves elements of EF (Bigorra et al., 2016). However, there were little or no far-effects on wider behavioural symptoms of ADHD in children. This may be because EF is only one of several endophenotypes of ADHD (Sonuga-Barke et al., 2014), EF deficits are not consistently represented across ADHD populations (Barkley & Murphy, 2010), and because ADHD is a heterogenous condition with many manifestations (Wahlstedt et al., 2009). Therefore, a multi-faceted, multi-modal approach may be more effective at reducing the behavioural symptoms of ADHD in children (Murray et al., 2008).

Including CMWT in a multi-modal approach raises the ethical issue of cost-verses-benefit (Guyatt et al., 2000), which would not work in CWMT's favour, as the costs are high. Furthermore, one of the selling points of CWMT is its usability in the home, reducing the need for on-site professional input. Chacko et al. (2014) raised leatteno ohefr (i)6 2 ( )10nssiing then (Mani2 ( )10nsee

suggestions for consideration are motivational or energetic endophenotypes (Sonuga-Barke, 2014). These interventions may prove more successful, especially if they are used synergistically to reduce the behavioural symptoms of children with ADHD.

5.0. References

American Psychiatric Association. (1994). *Diagnostic and Statistical Manual of Medical Disorders* (4th ed.). APA.

CAMHS Advisory Group (2018). *Delivering Effective Services for Children and Young People with ADHD*. CAMHS Advisory Group. https://www.england.nhs.uk/north-west/wp-content/uploads/sites/48/2019/03/GM-wide-ADHD-guidance.pdf

Castellanos, F. X., & Tannock, R. (2002). Neuroscience of attention-deficit/hyperactivity disorder: The search for endophenotypes. *Neuroscience, 3*, 617-628. https://doi.org/10.1038/NRN896

Cortese, S., Ferrin, M., Brandeis, D., Buitelaar, J., Daley, D., Dittman, R. W., Holtmann, M., Santosh, P., Stevenson, J., Stringaris, A., Zuddas, A., & Sonuga-Barke, E. J. S. (2015). Cognitive training for attention-deficit/hyperactivity disorder: Meta-analysis of clinical and neuropsychological outcomes from randomized controlled trials.

,0(eM)7(e2)12(16)3(aa)10 (r)20 0(r)4(i)7(a)1(3)7(Ri)6 um[hs.44 53(s)12)Ris,&(r)77 al 14

Guyatt, G. H., Haynes, R. B., Jaeschke, R. Z., Cook, D. J., Green, L., Naylor, C. D., Wilson, M. C., & Richardson, W. S. (2000). Users' guides to the medical literature XXV. Evidence-based medicine: Principles for applying the users' guides to patient care. *Journal of American Medical Association, 284*(10), 1290-1296. https://doi.org/10.1001/jama.284.10.1290

Guyatt, G. H., Sackett, D. L., Sinclair, J. C., Hayward, R., Cook, D. J., & Cook, R. J. (1995). Users' guides to the medical literature IX. A method for grading health care recommendations. *Journal of American Medical Association, 274*(22), 1800-1804.

Gwernan-Jones, R., Moore, D. A., Cooper, P., Russell, A. E., Richardson, M., Rogers, M., Thompson-Coon, J., Stein, K., Ford, T. J., & Garside, R. (2016). A systematic review and synthesis of qualitative research: the influence of school context on symptoms of attention deficit hyperactivity disorder. *Emotional and Behavioural Difficulties, 21*(1), 83-100. https://doi.org/10.1080/13632752.2015.1120055

Hague, C., Holland, A. A., & Gottleib, M. (2020). Selected ethical issues in computerized cognitive training. *Clinical Practice in Pediatric Psychology, 8*(4), 402-408. https://doi.org/10.1037/cpp0000311

evidence. *Child Adolescent Psychiatric Clinics of North America, 23*, 807-824. http://dx.doi.org/10.1016/j.chc.2014.05.009

Steiner, N. J., Frenette, E. C., Rene, K. M., Brennan, R. T., & Perrin, E. C. (2014). In-school neurofeedback training for ADHD: Sustained improvements from a randomized control trial. *Pediatrics, 133*(3), 483-492. http://dx.doi.org/10.1542/peds.2013-2059

Taylor, E., Dopfner, M., Sergeant, J., Asherson, P., Banaschewski, T., Buitelaar, J., Coghill, D., Danckaerts, M., Rothenberger, A., Sonuga-Barke, E., Steinhausen, H-C, & Zuddas, A. (2004). European clinical guidelines for hyperkinetic disorder – first upgrade. *European Child and Adolescent Psychiatry, 13*(1), 7-30. https://doi.org/10.1007/s00787-004-1002-x

Van de Loos-Neus, G. H. H., Rommelse, N., & Buitelaar, J. K. (2011). To stop or not to stop? How long should medication treatment of attention-deficit hyperactivity disorder be extended? *European Neuropsychopharmacology, 21*, 584-599. https://doi.org/10.1016/j.euroneuro.2011.03.008

van Dongen-Boomsma, M., Vollebregt, M. A., Buitelaar, J. K., & Slaats-Willemse, D. (2014). Working memory training in young children with ADHD: A randomized placebo-controlled trial. *Journal of Child Psychology and Psychiatry, 55*(8), 886-896. https://doi.org/10.1111/jcpp.12218

Wahlstedt, C., Thorell, L. B., & Bohlin, G. (2009). Heterogeneity in ADHD: Neuropsychological pathways, comorbidity and symptom domains.

Journal of Abnormal Child Psychology, 37, 551-564. http://dx.doi.org/10.1007/s10802-008-9286-9

Wilson, D. B. (2022, July 20). Practical Meta-Analysis Effect Size Calculator [Online calculator]. Campbell Collaboration. https://campbellcollaboration.org/research-resources/effect-size-calculator.html

Zhang, S., Faries, D. E., Vowles, M., & Michelson, D. (2005). ADHD rating scale IV: Psychometric properties from a multinational study as a clinician-administered instrument. *International Journal of Methods in Psychiatric Research, 14*, 186-201. http://dx.doi.org/10.1002/mpr.7

6.0. Appendices

Appendix A: Studies Excluded at Full Text Screening with Rationale

Table A

To Show Studies Excluded at the Full Text Screening Stage of the Literature Search

Referenced Article	Exclusion Criteria	Rationale
Bikic et al. (2017)	Feasibility study – not a fully-fledged RCT. Only include RCTs.	Feasibility studies are underpowered and explore if a full RCT is justified (Abbott, 2014).
Klingberg et al. (2002).	Not using commercial software.	Difficult to establish the reliability of the software. Published software can be reviewed and the study replicated.
Prins et al. (2011).	Not using commercial software.	Difficult to establish the reliability of the software. Published software can be reviewed and the study replicated.
Wolte(y)14ID 47 e3(		

Bikic, A., Christensen, T, O., Leckman, J. F., Bilenberg, N., & Dalsgaard, S. (2017). A double-blind randomized pilot trial comparing computerized cognitive exercises to Tetris in adolescents with attention-deficit/hyperactivity disorder. *Nordic Journal of Psychiatry, 71*(6), 455-464. https://doi.org/10.1080/08039488.2017.1328070

Prins, P. J. M., Dovis, S., Ponsioen, A. P., ten Brink, E., & van der Oord, S. (2011). Does computerised working memory training with game elements enhance motivation and training efficacy in children with ADHD? *Cyberpsychology, Behaviour and Social Networking, 14*(3), 115-122. https://doi.org/10.1089/cyber.2009.0206

Woltering, S., Gu, C., Liu, Z-X., & Tannock, R. (2021). Visu6p2kpad tial

Appendix B: WoE A Adaptations to the Kratochwill (2003) Coding Protocol

Weight of Evidence (WoE) ratings were based on the Harden & Gough (2012) framework which provides systematic and objective ways to evaluate studies. WoE A assesses the Methodological Quality of a study. This was achieved using an adapted version of the Procedural Manual of the Task Force on Evidence-Based Interventions in School Psychology, American Psychology Association (Kratochwill, 2003). The Kratochwill (2003) protocol was chosen to assess WoE A because it was a robust protocol suited to the design of RCTs and all of the studies reviewed were RCTs.

Table B

To Show how the Kratochwill (2003) Coding Protocol was Adapted for the Review

Section of Protocol Excluded	Rationale for Exclusion
Domain	This was reduced to 'School- and community-based intervention programmes for social and behavioural problems' and the other options deleted. This was because the intervention was for social and behavioural problems and the inclusion criteria stated that interventions should be practical enough for use at school and home. Thus, this box would be ticked for every study reviewed.
Part I General Study Characteristics	
A1 to A5	The study characteristics were described in the Mapping the Field table (Table 5), so this was considered duplication.
C3 Effect Size	

Section of Protocol Excluded	Rationale for Exclusion
H1.4, H2, H3, H4 Follow-up	Excluded as not considered to add to the findings. Information in H1.1 to H1.3 was considered sufficient to discriminate between studies.
I Identifiable intervention components	Excluded - not considered relevant as the intervention was a very specific computer-based working memory training programme and all studies used the same programme.
J1 Intervention Fidelity	Not considered it would add to differentiating between studies as J2 – manualisation – gave sufficient information.
J4.1 to J4.4 Implementation Context	These aspects were not considered relevant to the review and would be difficult to assess.
J4.7 Implementation Context Dosage response	Not considered relevant as dosage was not part of the study.
J4.9 Intervention style	Not considered to add to the review as the same computer programme was used for the intervention in all of the studies reviewed.
J4.10 Cost analysis data	Considered beyond the remit of this study.

Appendix C: WoE A Ratings for Methodological Quality

Weight of Evidence A Ratings

The WoE A judgements for each study reviewed were summarised in Table C.

The scores represented the following judgements:

- 3 = Strong Evidence
- 2 = Promising Evidence
- 1 = Weak Evidence
- 0 = No/Limited Evidence.

The scores for each indicator represented the average score of the sub-sections of each indicator, hence the decimalised scores.

The following key was used to describe the average scores calculated:

Key:

Range used to assign descriptors to mean scores:

Appendix D: WoE A Scoring Criteria

The scores in Table C were assigned according to criteria for each indicator. The criteria are described in Table D.

Table D

Summarising the criteria used to assign scores to assess the studies for WoE

A.

	Criteria used to assign scores of …

Appendix E: WoE B Ratings for Methodological Relevance

WoE B focused on:

1. Data Analysis: Is the sample size adequately powered to justify statistical analyses?
2. External Validity Indicators: Were the strategies employed appropriate to recruit participants with ADHD?
3. Implementation Fidelity: Was adequate support provided for implementers to deliver the intervention effectively?

Key:

Range used to assign descriptors to mean scores:

0.00 – 1.09 = No/Limited Evidence

1.10 – 1.69 = Weak

1.70 – 2.29 = Promising

2.30 – 3.0

Table E

To show a summary of WoE B scores for Methodological Relevance

Study	Criterion 1: Statistical Power	Criterion 2: Recruitment Strategy	Criterion 3: Support for Implementers	Overall WoE B	Descriptor
1. Bigorra et al. (2016)	3	3	1	2.33	Strong
2. Chacko et al. (2014)	1	1	2	1.33	Weak
3. Green et al. (2012)	0	2	2	1.33	Weak
4. Hovic et al. (2013)	0	3	3	2.00	Promising
5. Klingberg et al. (2005)	0	2	1	1.00	No/Limited Evidence
6. Van Dongen-Boomsma et al. (2014)	2	2	1	1.67	Weak
Mean	1.00	2.16	1.67	1.61	
Descriptor	No/Limited Evidence	Promising	Weak	Weak	

**NB: A score of zero indicates that one or none of the criteria is met or there is insufficient evidence provided.*

Table F

To show WoE B criteria for Methodological Relevance

	Criteria	*Ratings	Rationale
1	Is the sample size		

Appendix F: WoE C Ratings for Study Relevance to the Review Question

WoE C focused on:

1. Follow-Up: Does the study conclude whether or not CWMT improves behavioural symptoms of children with ADHD?
2. Measurement: Did the primary outcome measures effectively measure children's behavioural ADHD symptoms?
3. External Validity Indicators: Was children's ADHD status confirmed effectively?

Key:

Range used to assign descriptors to mean scores:

0.00 – 1.09 = No/Limited Evidence

1.10 – 1.69 = Weak

1.70 – 2.29 = Promising

2.30 – 3.00 = Strong

Table H

To show WoE C criteria for Study Relevance to the Review Question

	Criteria	*Ratings	Rationale
1	Does the study conclude whether or not CWMT improves the behavioural symptoms of children with ADHD?	3. Most primary outcomes, including follow-up outcomes, align, confirming whether or not CWMT improves the behavioural symptoms of ADHD in children. 2. Most outcome measures align, confirming whether or not CWMT improves the behavioural symptoms of ADHD in children, but there is no follow-up data. 1. Outcome measures are inconsistent such that whether or not CWMT improves the behavioural symptoms of ADHD in children with ADHD is inconclusive.	The main challenge for children with ADHD is conforming to the behavioural expectations of others. Therefore, it is important that CWMT improves children's behaviour, and that this is a key outcome for consideration when assessing the impact of CWMT on the behavioural symptoms of children with ADHD.
2	Did the primary outcome measures effectively measure children's behavioural ADHD symptoms?	3. At least two primary outcome measures measure a behavioural symptom of ADHD in children, with reliability data. 2. At least two primary outcome measures measure a behavioural symptom of ADHD in children. 1. At least one primary outcome measure measures a behavioural symptom of ADHD in children.	In order to assess if CWMT improves the behaviour of children with ADHD, it is importantavi behhavi3.12sr (s)/ in child measDHD, it is im r-i0 (e)]TJT*[(b[(thhione-6 (r-04 Tc 2

	Criteria	*Ratings	Rationale
3	Was children's ADHD status confirmed effectively?	3. ADHD status confirmed by clinical assessment prior to screening. 2. ADHD status confirmed by clinical assessment during screening. 1. Non-clinical screening for ADHD status.	The ADHD status of child

Appendix H: WoE A Coding Protocols

[Adapted from the Procedural Manual of the Task Force on Evidence-Based Interventions in School Psychology, American Psychology Association, Kratochwill, T.R. (2003)]

Coding Protocol: Group-Based Design

Domain: School- and community-based interventions for social and behavioural problems.

Name of Coder: TEP	**Date:** 17/06/2022
Full Study Reference in APA format: Bigorra, A., Garolera, M., Guijarro, S., & Hervas, A. (2016). Long-term far-transfer effects of working memory training in children with ADHD: A randomised controlled trial. *European Child and Adolescent Psychiatry, 25,* 853-867.	
Intervention Name (from description of study): Working memory computer training	
Study ID Number (Unique identifier): 01	

Type of Publication: Journal article

Part I

A. General Design Characteristics

A1. Random assignment designs (if random assignment design, select one of the following)

☒ Completely randomized design.
☐ Randomized block design (between participants, e.g., matched classrooms).
☐ Randomized block design (within participants).
☐ Randomized hierarchical design (nested treatments).

☐ 1 – Weak Evidence ☐ 0 - No Evidence.

B. Data Analysis

B1. Appropriate unit of analysis? Yes No

☐ 2 Vague or no links established between research methods and sampling, but sampling is *appropriate* to the research methods.
☐ 1 Links established between research method and sampling, but sampling is *inappropriate* to the research methods.
☐ 0 No links are established and sampling is inappropriate to research methods.

A2. Operationalisation. Specifying the link between key abstract constructs (variables)

☒ 3 Clear links established between constructs and methods, **and** all key constructs are clearly operationalised.
☐ 2 Some, but not all, key constructs are clearly operationalised.
☐ 1 Vague reference to link between constructs and methods.
☐ 0 No evidence that key constructs are operationalised.

A3. Integration of data from multiple sources, methods and investigators.

☒ 3 Used multiple sources, methods, and investigations.
☐ 2 Used two of the following: multiple sources, multiple methods, multiple investigators.
☐

B4. Extent of engagement. The researchers conduct data collection in a manner that guarantees sufficient scope and depth through prolonged engagement (data collection over a sufficient time period to ensure accuracy of representation) and persistent observation (progressively focused to ensure thorough understanding of consistency and variation), respectively.

☐ 3 Provided evidence for high level of engagement to ensure deep and accurate representation.
☒ 2 Provided evidence for some level of engagement to ensure deep and accurate representation.
☐ 1 Provided minimal l9 (h l)6 (ev)1-0.004 Tw 4.45 0 Td[(epr)7 (es)14 (ent)12 (at)2 (i)6 (o)

☒ 3 – Strong Evidence ☐ 2 – Promising Evidence
☐ 1 – Weak Evidence ☐ 0 - No Evidence.

D. External Validity Indicators

D1. Sampling procedures

D1.1 Sampling procedures described in detail

☒ Yes ☐ No (incomplete or no evidence)

D1.2. Rationale for sample selection specified

☒ Yes ☐ No (incomplete or no evidence)

D1.3. Rationale for sample size specified

☒ Yes ☐ No (incomplete or no evidence)

D1.4. Evidence that sample represents target population

☒ Yes ☐ No (incomplete or no evidence)

D1.5. Recruitment procedures congruent with target cultural group. Researcher used culturally appropriate ways/methods to contact, recruit, inform and maintain participation.

☒ Yes ☐ No (inadequate description or no evidence)

D1.6. Inclusion/exclusion criteria specified ☒ Yes ☐ No

D1.7. Specified criteria related to concern ☒ Yes ☐ No

D1. Overall rating on Sampling Procedures

☒ 3 – Strong Evidence ☐ 2 – Promising Evidence
☐ 1 – Weak Evidence ☐ 0 - No Evidence.

D2. Participant Characteristics specified for treatment and control groups (select all that apply).

☐

☒ 3 Complete and detailed description of the context within which the intervention occurs.
☐ 2 Detailed description of some but not all contextual components.
☐ 1 Provides overview of contextual components but lack details.
☐ 0 No description of context.

D2 and D3 Rating for Participant Characteristics and Transferability of the Intervention

☒ 3 – Strong Evidence ☐ 2 – Promising Evidence
☐ 1 – Weak Evidence ☐ 0 - No Evidence.

E. Follow-up Assessment

E1. Timing of follow-up assessment

☒ Yes ☐ No ☐ Unclear

Specify _1-2 weeks post-training and 6 months post-training

E2. Number of participants included in the follow up assessment

☒ Yes ☐ No ☐ Unclear

Specify __55/66.

F1.3 Written material involving an overview of broad principles and a description of the intervention phases.

F1.4 Formal or informal training session involving an overview of broad principles and a description of the intervention phases.

F2 Adaptation procedures are specified ☐ Yes ☐ No ☒ Unknown

F1. and F2. Rating for Manualisation and Adaptation

☐ 3 – Strong Evidence
☒ 1 – Weak Evidence
☐ 2 – Promising Evidence
☐ 0 - No Evidence.

F3 Conditions of intervention

F3.1. Length of intervention

Unknown/Insufficient information provided

Information provided (if information provided, specify the following:)

☐ Weeks _5
☐ Months _____
☐ Years _____
☐ Other _____

F3.2. Intensity/dosage of intervention

☐ Frequency of intervention session _5 sessions per week (25 total)
☐ Length of intervention session _35-45mins/ 90 trials
☐ Unknown/insufficient information provided _____

F3.1 and F3.2. Rating length of intervention and intensity

☐

Summary of Evidence