developmentally appropriate commands in an authoritative parenting style and must follow through with consequences for compliance and non-compliance (Herschell et al., 2002).If the child complies then the adult must give specific praise and if the child does not comply then a time-out will be initiated (Herschell et al., 2002). The intervention takes between 10 and 16 weeks to complete usually with 1-hour

as they are able to immediately recognise and change 'established patterns of verbalization and behaviour with their child' (Eyberg, 1988).

Attachment theory suggests that care-givers and children can form different attachment types based on the responsiveness of the care-giver to child which can predict behaviours for children in the future. A secure attachment type is formed when caregivers respond to a child's needs regularly and appropriately and this helps children to form a positive schema of relationships, good emotional regulation and good social skills (Ainsworth, 1979). On the other hand, when a caregiver is less responsive or inconsistent, they can form an insecure attachment with the child which is a risk factor for children to develop behavioural difficulties (Ainsworth, 1979). The skills parents learn in the child-directed interaction phase fosters and promotes a secure attachment with the child which increases positive interaction and responsiveness between parent and child (Serchuk et al., 2021). This is why the parents are encouraged first of all to engage in child-interacted play to increase a positive bond (Serchuk et al., 2021). The PDI phase is influenced by social learning theory, developmental theory and behavioural theory with the aim to build behaviour management skills for the caregiver and strategies to help the child learn, through modelling, desired behaviours (Serchuk et al., 2021).

Autism

Autism spectrum condition (ASC), also known as autism spectrum disorder (ASD) is a neurological difference that impacts three main characteristics of a person (Faras et al., 2010). Differences are usually noticed in communication, social interaction and repetitive patterns of behaviour which can present with issues with adapting to change, making positive friendships, ability to focus and problem solving (Faras et al., 2010). Parents and schools have reported significant challenges with

used to measure

the literature search process. Included in the final review was 7 studies, in Table 3 they are summarised but further detail can be found described in Appendix B.

Table 2

Rationale for exclusion and inclusion criteria:

Criteria	Inclusion criteria	Exclusion criteria	Rationale
1. Participants	x Must be between 2-17 years old x Must have diagnosis of autism	x Over 18 years old x Dual diagnosis for example ASD and ADHD	x We want to find out effectiveness of intervention for children who have autism
2. Settings	x Intervention carried out in community, professional work place or at care-givers home	x Controlled setting	x Maintain fidelity of intervention
3.			

Figure 1 :

PRISMA flow chart showing inclusion and exclusion process:

Table 3

A summary of final studies included in the review.

No.	Studies included
1.	Han, R. C., Naguib, S., Owen, C. K., Druskin, L. R., Keen, K. R., Piper, R., Holbert, S. N., Shank, S. D., Victory, E. J., & McNeil, C. B. (2022). An Effectiveness Trial of PCIT for Children with and without Autism Spectrum Disorder in a Private Practice Setting.

Adolescent Mental Health, 7(1), 125–141. Child Development &

Adolescent Studies. https://doi.org/10.1080/23794925.2021.1993109

2. Zlomke, K. Q 0 g 0002 Tw 12 -0 0M1 ( )]TJ( )J (S)1(ps)4e(/)2 ,TJ E92 (Q 0

6. Solomon, M., Ono, M., Timmer, S., & Goodlin-jones, B. (2008). The Effectiveness of Parent-Child Interaction Therapy for Families of Children on the Autism Spectrum. Journal of Autism and Developmental Disorders, 38(9), 1767–1776. ProQuest Central; Social Science Premium Collection. https://doi.org/10.1007/s10803-008-0567-5

7. Quetsch, L. B., Bradley, R. S., Theodorou, L., Newton, K., & McNeil, C. B. (2022). Community-based agency delivery of Tw 1014 (or)-hm (7S)1 (.)12 . ET Q .Chileol2 onol7tifputism durna(on1pm)17 (ent a()7 (2d.)2)1 (p(on)10d(A)11 (ut)2 (i)14 (m)-3 ((enes)4)

Overall weight of evidence

Study	Ratings				Rating descriptor
	WoE A	WoE B	WoE C	WoE D- overall rating	
Han et al., 2022	2.75	2	3	2.6	High
Zlomke, &; Jeter., 2020	2.75	2	3	2.6	High
Scudder et al., 2019	2.5	3	3	2.8	High

research into longevity of its impact. Attrition

Measurement

The Eyberg child behaviour inventory (Eyberg & Pincus, 1999) was used to measure the frequency of negative behaviour in the participants. It is used by care-givers to report on the disruptive behaviours of their children through a 7-point Likert scale for subscale Intensity and a “yes” or “no” scale for subscale Problem. Data from this measure was collected at pre, mid and post intervention to see the effect of the intervention. Most studies reported the inter-reliability coefficient for ECBI and all studies reported on the scale’s reliability from different sources. The weight of reliability of this measurement was recorded in WoE A in the feature ‘Measurement’ which impacted the consequent score. Each study also measures parent outcomes however this is not considered and it is not the focus of this review. Other measurements that observed child outcomes were also used and reported in Mapping of the Field in Appendix B however, only ECBI was considered in this review as it was used as a primary measurement in all studies.

Outcomes and Effect sizes

Han et al (2022) revealed promising outcomes for the effectiveness of PCIT on children with autism]. However did not have enough sample size power to detect a statistically significant change as there was a difference in pre to post scores for ECBI; the effect sizes were small according to Cohens d as reported in Table 5. Due to COVID-19 the authors were not able to recruit more participants to increase the power of the study. A statistically significant effect was found for both groups from pre to post treatment on both subscales of the ECBI in the Quetsch et al (2022) study although small and medium effect sizes were found as reported in Table 5. This suggests a PCIT is effective at reducing negative behaviours in children with autism.

All other studies showed a statistically significant main effect for time for the group of children with a. This suggests the intervention had an impact before it was delivered and after which shows strong evidence that PCIT can be effective at reducing negative behaviours of children with autism. Most studies looking at the efficacy of PCIT had comparison groups, including wait-list control has shown that the treatment condition is more effective than control settings.(Herschell et al., 2002). Solomon et al (2008) yielded a significant main effect for group X time for the problem scale which shows there is a difference between groups of how parents perceived problematic nature of their behaviours. As this study design delivers a wait-list control, this shows promising evidence for PCIT having positive effects on parent-led interventions impacting the behaviours of children with autism. All studies, apart from Han et al (2022) and Quetsch et al (2022) revealed large to huge effect sizes shown in Table 5. Cohens d (Cohen, 1988) was used to interpret effect sizes for majority of studies. Quetsch et al (2022) reported partial eta squared as shown in the effect size tables.

Table 5

Table showing effect sizes and WoE overall rating

Study	Sample size	Groups comparison	Outcome measurements	Effect size type	Effect size (ES)	ES descriptor [a]	Overall Weighting (WoE D)
(Han et al., 2022)	22	ASD vs non-ASD	ECBI	Cohens d	Problem = 0.09 Intensity= 0.18	Small Small	High
(Zlomke & Jeter ., 2020)	28	ASD vs non-ASD	ECBI	Cohens d	Intensity= 2.27	High	High

[a] Cohen' d (1988) interpreted as d = .10 (small), d = .50 (medium), d = .80 (large) and d = .1.2 (very large) and d = 2.0 (Sawilosky, 2009)
*Partial eta squared interpreted as $_{p}^{2}$ = .01 (small), $_{p}^{2}$ = .06 (medium) and $_{p}^{2}$ = .14 (large) (Cohen et al., 2014)

References:

Ainsworth, M. S. (1979). Infant–mother attachment. American psychologist, 34(10), 932.

Agazzi, H., Tan, S. Y., Ogg, J., Armstrong, K., & Kirby, R. S. (2017). Does parent-child interaction therapy reduce maternal stress, anxiety, and depression among mothers of children with autism spectrum disorder? Child & Family Behavior Therapy, 39(4), 283–303. . https://doi.org/10.1080/07317107.2017.1375622

Allen, K., Harrington, J., Quetsch, L. B., Masse, J., Cooke, C., & Paulson, J. F. (2023). Parent–Child Interaction Therapy for Children with Disruptive Behaviors and Autism: A Randomized Clinical Trial. Journal of Autism and Developmental Disorders, 53(1), 390–404. https://doi.org/10.1007/s10803-022-05428-y

Baumrind, D. (1991). The Influence of Parenting Style on Adolescent Competence and Substance Use. The Journal of Early Adolescence, 11(1), 56–95. https://doi.org/10.1177/0272431691111004

Bradshaw, J., Steiner, A. M., Gengoux, G., & Koegel, L. K. (2015). Feasibility and Effectiveness of Very Early Intervention for Infants At-Risk for Autism Spectrum Disorder: A Systematic Review. Journal of Autism and Developmental Disorders, 45(3), 778–794. https://doi.org/10.1007/s10803-014-2235-2

Cohen, J. (1988). Set Correlation and Contingency Tables. Applied Psychological Measurement, 12(4), 425–434. https://doi.org/10.1177/014662168801200410

Cohen, P., Cohen, P., West, S. G., & Aiken, L. S. (2014). Applied Multiple Regression/Correlation Analysis for the Behavioral Sciences (2nd ed.). Psychology Press. https://doi.org/10.4324/9781410606266

Eyberg, S. (1988). Parent-Child Interaction Therapy: Integration of Traditional and Behavioral Concerns. Child & Family Behavior Therapy, 10(1), 33–46. https://doi.org/10.1300/J019v10n01_04

Eyberg, S. M., & Pincus, D. (1999). Eyberg Child Behavior Inventory and Sutter-Eyberg Student Behavior Inventory: Professional manual. Odessa, FL: Psychological Assessment Resources.

Faras, H., Al Ateeqi, N., & Tidmarsh, L. (2010). Autism spectrum disorders. Annals of Saudi Medicine, 30(4), 295–300. https://doi.org/10.4103/0256-4947.65261

Fawley, K. D., Stokes, T. F., Rainear, C. A., Rossi, J. L., & Budd, K. S. (2020). Universal TCIT Improves Teacher–Child Interactions and Management of Child Behavior. Journal of Behavioral Education, 29, 635–656. https://doi.org/10.1007/s10864-019-09337-6

Gershenson, R. A., Lyon, A. R., & Budd, K. S. (2010). Promoting Positive Interactions in the Classroom: Adapting Parent-Child Interaction Therapy as a

Universal Prevention Program. Education and Treatment of Children, 22(2). https://doi.org/10.1353/etc.0.0092

Gough, D. (2007). Weight of Evidence: A framework for the appraisal of the quality and relevance of evidence. Special Edition of Research Papers in Education, 22(2), 213–228. https://doi.org/10.1080/02671520701296189

Han, R. C., Naguib, S., Owen, C. K., Druskin, L. R., Keen, K. R., Piper, R., Holbert, S. N., Shank, S. D., Victory, E. J., & McNeil, C. B. (2022). An Effectiveness Trial of PCIT for Children with and without Autism Spectrum Disorder in a Private Practice Setting. Evidence-Based Practice in Child & Adolescent Mental Health, 7(1), 125–141. Child Development & Adolescent Studies. https://doi.org/10.1080/23794925.2021.1993109

Hanf, C. (1968). Modifying Problem Behaviors in Mother-Child Interaction: The Standardized Laboratory Situations (64110999; ED087561). 1–23. ERIC. https://www.proquest.com/scholarly-journals/modifying-problem-behaviors-mother-child/docview/64110999/se-2?accountid=14511

Herschell, A. D., Calzada, E. J., Eyberg, S. M., & McNeil, C. B. (2002). Parent-child interaction therapy: New directions in research. Cognitive and Behavioral Practice, 9(1), 9–16. https://doi.org/10.1016/S1077-7229(02)80034-7

Kratochwill, T. R. (2003). TASK FORCE ON EVIDENCE-BASED INTERVENTIONS IN SCHOOL PSYCHOLOGY.

Lesack, R., Bearss, K., Celano, M., & Sharp, W. G. (2014). Parent-Child Interaction Therapy and autism spectrum disorder: Adaptations with a child with severe developmental delays. Clinical Practice in Pediatric Psychology, 2(1), 68–82. https://doi.org/10.1037/cpp0000047

Masse, J. J., McNeil, C. B., Wagner, S. M., & Chorney, D. B. (2007). Parent-child interaction therapy and high functioning autism: A conceptual overview. Journal of Early and Intensive Behavior Intervention, 4(4), 714–735. APA PsycInfo <2006 to 2007>. https://doi.org/10.1037/h0100402

Masse, J. J., Mcneil, C. B., Wagner, S., & Quetsch, L. B. (2016). Examining the Efficacy of Parent-Child Interaction Therapy with Children on the Autism Spectrum. Journal of Child and Family Studies, 25(8), 2508–2525. ProQuest Central; Social Science Premium Collection. https://doi.org/10.1007/s10826-016-0424-7

Matano, M., Monden, Y., Kurane, K., Kawasaki, M., & Kamo, T. (2021). Potential of internet-delivered PCIT for ASD in the COVID-19 era: A pilot study. Pediatrics International. Coronavirus Research Database. https://doi.org/10.1111/ped.14699

McInnis, P., Kohlhoff, J., & Eapen Valsamma. (2020). Real-world Outcomes of PCIT for Children at Risk of Autism or Developmental Delay. Journal of Child and Family Studies, 29(6), 1701–1711. ProQuest Central; Social Science Premium Collection. https://doi.org/10.1007/s10826-020-01699-0

Solomon, M., Ono, M., Timmer, S., & Goodlin-jones, B. (2008). The Effectiveness of Parent-Child Interaction Therapy for Families of Children on the Autism Spectrum. Journal of Autism and Developmental Disorders, 38(9), 1767–1776. ProQuest Central; Social Science Premium Collection. https://doi.org/10.1007/s10803-008-0567-5

Stichter, J. P., Riley-Tillman, T. C., & Jimerson, S. R. (20161208). Assessing, understanding, and supporting students with autism at school: Contemporary science, practice, and policy. School Psychology Quarterly, 31(4), 443. https://doi.org/10.1037/spq0000184

Vess, S. F., & Campbell, J. M. (2022). Parent–child interaction therapy (PCIT) with families of children with autism spectrum disorder. Autism & Developmental Language Impairments, 7. ProQuest Central; Social Science Premium Collection. https://doi.org/10.1177/23969415221140707

Zlomke, K. R., & Jeter, K. (2020). Comparative Effectiveness of Parent–Child Interaction Therapy for Children with and Without Autism Spectrum Disorder. Journal of Autism and Developmental Disorders, 50(6), 2041–2052. ProQuest Central; Social Science Premium Collection. https://doi.org/10.1007/s10803-019-03960-y

Zlomke, K. R., PhD, Jeter, K., MS, & Murphy, J., MS. (2017). Open-Trial Pilot of Parent-Child Interaction Therapy for Children With Autism Spectrum Disorder. Child & Family Behavior Therapy, 39(1), 1–18. ProQuest Central; Social Science Premium Collection. https://doi.org/10.1080/07317107.2016.1267999

Lesack, R., Bearss, K., Celano, M., & Sharp, W. G. (2014). Parent–Child Interaction Therapy and autism spectrum disorder: Adaptations with a child with severe developmental delays. Clinical Practice in Pediatric Psychology, 2(1), 68. https://doi.org/10.1037/cpp0000047	Study design: Study did not have a control group design and did not have pre and post conditions measures of child outcomes
Masse, J. J., McNeil, C. B., Wagner, S. M., & Chorney, D. B. (2007). Parent-child interaction therapy and high functioning autism: A conceptual overview. Journal of Early and Intensive Behavior Intervention, 4(4), 714. https://doi.org/10.1037/h0100402 (Masse et al., 2007)	Study design: Study did not have a control group. Used case study design

Appendix C Weight of Evidence A

In this review the APA Kratochwill protocol (2003) was used to weigh the methodology of each study. Below in table 1 you will find the rationale for modifications to the protocol, as only what was relevant to the weight of Evidence A was used.

Table 1

Amendments made to Kratochwill (2003) Coding protocol

Sections	Sections removed /modified	Rationale
General Study Characteristics	A1 – A5	Discussed in literature review
Data Analysis	Section modified C1-C3 C7-C9	Data described in literature review and some information is reporting in the Mapping the field table. Secondary outcomes are not relevant to review question.
Research Methodology	Whole section removed	Stated in Mapping of Fields table
Measurement	Section modified B4 – B7	Data added within literature review
Primary/Secondary Outcomes are	D1 – D6	Primary outcomes are discussed in literature review. Secondary outcomes are not applicable to

		review question as children outcomes are being looked at
Statistically Significance	E, F, G & H	Discussed in literature review and section H not relevant to review of methodological quality
Implementation Fidelity	J4.1-J4.12	Review of an evidence-based intervention that has a manualised approach
Implementation fidelity context	K & L	Review of an evidence-based intervention that has a manualised approach that must be implemented by a qualified practitioner trained in the intervention. Discussed in literature review.

Appendix D: Kratochwill (2003) Coding protocol Weight of Evidence A

Adapted from the Procedural Manual of the Task Force on Evidence-Based Interventions in School Psychology, American Psychology Association, Kratochwill, T.R. (2003)]

Coding<</Mei 37 (a)1 (n5 (.)-8 (R)-4 (.)Me)5 (i 37 2illia)5 (o-0 0

A1. Random assignment designs (if random assignment design, select one of the following)

☐ Completely randomized design

☐ Randomized block design (between participants, e.g., matched classrooms)

☐ Randomized block design (within participants)

☐ Randomized hierarchical design (nested treatments

A2. Nonrandomized designs (if non-

Intervention group sample size: ___11____

Control group sample size: ____11____

C. Type of Program

☐ Universal prevention program
☐ Selective prevention program
☐ Targeted prevention program
☒ Intervention/Treatment
☐ Unknown

D. Stage of Program

☐ Model/demonstration programs
☐ Early stage programs
☒ Established/institutionalized programs
☐ Unknown

E. Concurrent or Historical Intervention Exposure

☒ Current exposure
☐ Prior exposure
☐ Unknown

2. Key Features for Coding Studies and Rating Level of Evidence/Support

(Rating Scale : 3= Strong Evidence, 2=Promising Evidence, 1=Weak Evidence, 0=No Evidence)

A. Measurement (Estimating the quality of the measures used to establish effects)

A1 The use of the outcome measures produce reliable scores for the majority of the primary outcomes

☒Yes

☐No

☐Unknown/unable to code

A2 Multi-method (at least two assessment methods used)

☒ Yes

☐ No

☐ N/A

☐ Unknown/unable to code

A3 Multi-source (at least two sources used self-reports, teachers etc.)

☒ Yes

☐ No

☐ N/A

☐ Unknown/unable to code

A4 Validity of measures reported (well-known or standardized or norm-referenced are considered good, consider any cultural considerations)

☒ Yes validated with specific target group

☐ In part, validated for general population only

☐ No

☐ Unknown/unable to code

- ☐ By change agent
- ☐ Statistical (analyse includes a test for intervention)
- ☒ Other
- ☐ Not reported/None

B4 Group equivalence established (select one of the following)

- ☐ Random assignment
- ☒ Posthoc matched set
- ☐ Statistical matching
- ☐ Post hoc test for group equivalence

B5 Equivalent mortality

- ☒ Low attrition (less than 20 % for post)
- ☒ Low attrition (less than 30% for follow-up)
- ☐ Intent to intervene analysis carried out?h(al)6 (enc)4 (e)]TJ 0 Tc (ha)d [(Ih(al)6h(al)al)6h(

Overall rating for Statistical Analysis __2__

3= Strong Evidence 2=Promising Evidence 1=Weak Evidence 0=No Evidence

D. Implementation Fidelity

D1. Evidence of Acceptable Adherence (answer J1.1 through J1.3)

D1.1 ☐

Summary of Evidence

Indicator	Overall evidence rating 0-3	Description of evidence Strong Promising Weak No/limited evidence Or Descriptive ratings
General Characteristics		
Design		Promising
Type of programme		Strong
Stage of programme		Strong
Concurrent/ historical intervention exposure		Promising
Key features		
Measurement	3	strong
Comparison group	3	Strong
Appropriate Statistical Analysis	2	promising
Implementation fidelity	3	Strong

Appendix E: Weight of Evidence A

- Below you will find the criteria, ratings and rationale for Weight of Evidence A, all of which has been used from Kratochwill (2003) protocol and adapted for relevance to this review question. Table 1.1 is the rationale and ratings for key features of each study
- Table 1.2 is the breakdown of ratings given to each study on one table
- Table 1.3 is the WoE A average rating scores

Table 1.1

WoE A: Rationale and ratings. Studies must meet all the criteria in each rating to achieve that ratings.

Key features	Rating

	- Measured outcomes must also reflect a moderate effect size
2 (Medium)	- statistical analysis must have been conducted with appropriate units of analysis - must show significant and primary outcomes for at least 50% to 74% of the total primary outcome measures
1 (Low)	-

Solomon et al .

Appendix F: Weight of Evidence B

In this section you will find the criteria, rational and ratings for Weight of Evidence B according to the hierarchy of evidence (Petticrew, 2003). This will give an indication of the appropriateness of study design used by each study (Gough, 2007).

- Table 1.1 is the evidence hierarchy of method design and their ratings according to Petticrew (2003)
- Table 1.2 is the criteria and rational used to gi gi8d0.002 Tw 1.19 071 (i)1

Table 1.3

WoE B: Overall Rating

Studies	Design method	WoE B rating