MCC 5th Conference



Daresbury Laboratory

Wednesday 28th June – Friday 30th June, 2023

Sponsors

The MCC is very grateful for funding from EPSRC.

We are grateful that this event is also sponsored by:

STFC, Daresbury Laboratory

Department of Chemistry (UCL)

Organising Committee

Georgia Lomas, Alin-Marin Elena, Thomas Keal, Alexey Sokol, and Scott Woodley

We also would like to thank all members, particularly colleagues who have helped us complete our recent EPSRC proposal during 2022, our theme leaders and all of you, including our invited guests, who have joined us in Daresbury in-person or online.

MCC 5th Conference - Programme

Wednesday 28th June			
12:30	**Registration Desk Open**		**Daresbury Laboratory**
12:30	**Lunch**		
Session 1:	Low Dimensional Materials	**Chair** Umberto Terranova	
13:45	**Start Welcome Address**		
13:50	**Invited:** Disorder?		Neil Allan Bristol
14:20	Predicting the		

Session 4:	Surfaces and Interfaces	**Theme Leader** Matthew Watkins
11:30	The effect of ceria surface composition on scavenging reactive oxygen species	Khoa Minh Ta Huddersfield (mol)
11:50	Modelling the effect of H in the formation, stabilisation and evolution of voids in polycrystalline Cu	Vasileios Fotopoulos UCL (shl)
12:10	Simulating Hydrogenation in Magnesium Through the Application of Machine Learning Models	

Friday 30th June			
Session 7:	Materials Discovery	**Chair** Marco Molinari	
9:00	Optimisation of decision sequences from selection of elements to chemical formulae of functional materials		A. Vasylenko Liverpool (dar)
9:20	Computational Modelling of Wide Band Gap Transparent Conducting Oxide Sb_2O_5		Ke Li UCL (dos)
9:40	A computational study characterising the intrinsic and extrinsic doping profile of ZnS		S. Aggarwal UCL (dos)
10:00			

Talks

Disorder?

Neil Allan*

* *School of Chemistry, University of Bristol*

TBA

Multiscale modelling of the electrical conductivity of carbon nanotube films

Thomas R. Durrant*, Yvelin Giret, David Z. Gao, Alexander L. Shluger

*Department of Physics and Astronomy, UCL e05-nano-**shl***

Carbon nanotubes (CNTs) are a fascinating low dimensional material with very high electrical conductivity. CNTs hold significant technological promise due to their extremely high electrical conductivities. This is because metallic CNTs behave as ballistic conductors, which enables highly energy efficient electrical devices to be constructed from them. In many practical applications, CNTs are deposited in a disorganized film. One such example application is the use of a CNT film as resistance changing element in non-volatile computer memory. In order to model the electrical conductivity of such films, models of the electrical conductivity of individual CNT junctions must be combined with larger-scale models of their geometric structure.

In this project, a multiscale model of the electrical conductivity of CNT films was developed through the following steps: **(a)** A large library of CNT-CNT junction structures was simulated using density functional tight binding (DFTB) in combination with the nonequi h. The intrinsic electrical conductivity of individual CNTs is high and the primary source of electrical resistance

ONETEP, with some exemplar applications thrown in

Chris Skylaris

University of S

Current Challenges in Catalytic Science

Graham Hutchings

University of Cardiff

TBA

Mechanism of NH_3 Synthesis on Fe_3Mo_3N

M. D. Higham*, Constantinos D. Zeinalipour-Yazdi, Justin S. J. Hargreaves, C. Richard A. Catlow

**Department of Chemistry, University College London* *e05-react-cat*

Plane-wave Density Functional Theory (DFT) techniques are applied to investigate two distinct reaction mechanisms for ammonia (NH_3) synthesis over a thin-film model Fe_3Mo_3N(111) surface, namely an associative Mars van Krevelen mechanism and the conventional, dissociative, Langmuir Hinshelwood mecha

The Electronic Structure and Catalytic Activity of a Ru3 Cluster Embedded on Nitrogen-doped Graphene

Yao Zhao*,

Computational investigation of NiOOH, from bulk to catalysis

Ruotao Yang*, Alexey A. Sokol, C. Richard A. Catlow

**Department of Chemistry, University College London*

*e05-react-**cat***

Mitigating environmental and ecological damage caused by human emissions are amongst the grand challenges of the 21st century. Therefore, the transition from conventional fossil energy to renewable energy, including solar and wind energy is an increasingly important topic. Power-to-gas (P2G), a process of converting renewable energy into hydrogen gas via electrolyser, is considered to be promising due to its carbon-free nature. However, to produce green hydrogen,

Amorphous ZnO: Structure, Stability and Charge Trapping

David Mora Fonz

HTU, Mexico

TBA

Ab-Initio Molecular Dynamics Simulations of Substoichiometric Amorphous Alumina

J. Strand*, A. Shluger

*Department of Physics and Astronomy, University College London

e05-bulk-**shl**

Oxygen vacancies play an important role in the application of oxides in technology. The notion of a vacancy is well defined in the case of crystals, where a defect is simply any break from crystallographic order. In amorphous materials, where there is no long-range order, defects are less well defined. Amorphous oxides are widely used in technology, yet there has not been a great deal of focus on this ambiguity. Often, vacancies are modelled by deleting oxygen atoms from a stoichiometric amorphous model, previously produced using the melt-and-quench technique, and then relaxing the system geometry - . However, this may not be the appropriate way to model substoichiometry in amorphous oxides. Here, we use an alternative approach where we produce substoichiometric amorphous oxide models by quenching from an oxygen-deficient melt - . We use ab-initio molecular dynamics (AIMD) with a hybrid DFT functional to investigate the structure of substoichiometric alumina. The models produced are periodic and pseudo-amorphous, with 357 atoms in the unit cell. We find that there are vacancy-like point defects in the amorphous oxide, characterized by undercoordinated Al ions and formation of Al-Al dimers. In subsequent simulations, we take post-quench vacancies and anneal them with ab-initio MD at 1000K. The resulting configurations are similar to those produced with the in-melt method, suggesting that use of ab-initio MD allows for a fuller exploration of configuration space and ther

Overall, the results suggest that, while vacancies may be ill-defined in amorphous oxides, oxygen deficiency leads to the formation of point-defects with similar properties to crystallographic vacancies. Further, it is possible that AIMD is necessary to find the most stable configurations of these defects.

Unravelling Hydrogen Bonding Behaviours and their Impact on Proton Conductivity in Supramolecular Cages

Lei Lei, Sanliang Ling*

*Advanced Materials Research Group, Faculty of Engineering, University of Nottingham

e05-bulk-**lin**

Proton-exchange membrane (PEM) fuel cells hold promise as a solution for achieving a zero-emission economy. Recently, molecular cages have emerged as potential PEM candidates for stable and efficient fuel cells through structural design. However, due to their complex structures, understanding the proton transport dynamics and mechanisms using experimental or simulation methods has proven challenging. In this study, we employed *ab initio* molecular dynamics simulations to investigate these complex systems over extended timescales, spanning hundreds of picoseconds, to unravel these intricate systems. Additionally, we developed efficient python code to analyse the vast trajectory data generated. Our results highlight a significant disparity in the interaction of water with two molecular cages composed of the same functional groups, attributed to their distinct geometries. Notably, Cage 1, characterized by a spacious cavity, exhibits an(en-US)>> B1 0 0 1 290.23 401.71 Tms4 82.224 478.9 519.19 reTtir-3( )-31((P)413( )-3

Investigation of unintentional aggregation of impurity-related defects within 4H-SiC.

N. Smith*, A. Shluger

Department of Physics and Astronomy, UCL* *e05-bulk-shl***

To create p-doped regions within 4H-SiC MOSFET devices, 4H-SiC is bombarded with high energy Al^+ ions from gaseous AlH_3, during ion implantation. However, recent experimental studies have found

Simulation of correlation driven metal-insulator transitions in real solids using deep neural networks.

The effect of ceria surface composition on scavenging reactive oxygen species

Khoa Minh Ta*, David J. Cooke, Lisa J. Gillie, Roger M. Phillips, Marco Molinari

*Department of Chemistry, University of Huddersfield *e05-surfin-**mol***

Cellular activities produce reactive oxygen species (ROS), such as hydrogen peroxide, superoxide anions, and hydroxyl radicals, which can accumulate and produce oxidative stress. This is linked to many degenerative diseases and cancer. Ceria nanoparticles (CeNPs) have been reported to perform enzyme mimetic activities to scavenge RO

Modelling the effect of H in the formation, stabilisation and evolution of voids in polycrystalline Cu

Vasileios Fotopoulos *, Alexander L. Shluger*

Physics and Astronomy, University College London *e05-surfin-**shl***

Cu is used in a wide variety of applications ranging from power generation to microelectronics.

Study of Hydroxyl Nests in Zeolites

Alec Desmoutier

* *University College London*

surfin-cat

Bulk and Surface Contributions to Ionisation Potentials of Metal Oxides

Xingfan Zhang*, Alexey A. Sokol, C. Richard A. Catlow

*Department of Chemistry, University College London e05-power-**sok***

Obtaining absolute band edge positions in solid materials remains an inherently complex task from both experiment and theory, and indeed there is considerable confusion in the literature. Our work shows clearly that the ionisation potential (IP) and electron affinity (EA) of metal oxides emerge as surface-related parameters, exhibiting high sensitivity to various factors, such as material morphologies and processing

Bismuth oxychalcogenide materials for thermoelectric applications

Joseph M. Flitcroft,* Adel Althubiani, Jonathan M. Skelton

*Department of Chemistry, University of Manchester *e05-power-ske*

In order to effectively combat climate change a wide range of approaches are required. One route is to increase the efficiency of energy-intensive process to reduce the associated CO_2

Controlling the heat transport in thermoelectric materials

Jonathan M. Skelton*

*Department of Chemistry, University of Manchester

The Baskerville Tier

Oxygen-functionalised graphene: structure, properties and potential for phosphate sensing

Negar Mansouriboroujeni, Xue Yong, Natalia Martsinovich*

Department of Chemistry, University of Sheffield *e05-enviro-**nat***

Graphene is an outstanding material thanks to its excellent electronic and optical properties. Graphene is also a promising sensor material, which ha

An Efficient Model for sp-Lone Pair Cations

Woongkyu Jee*, Scott M. Woodley, Alexey A. Sokol

University College London, Department of Chemistry

e05-

Optimisation of decision sequences from selection of elements to chemical formulae of functional materials

A. Vasylenko[1], D. Antypov[1], V. Gusev[1], G. Darling[1], M. S. Dyer[1], M. J. Rosseinsky[1]

[*]*Department of Chemistry, University of Liverpool*

Computational Modelling of Wide Band Gap Transparent Conducting Oxide Sb_2O_5

Ke Li[ab], Joe Willis[ab], David O. Scanlon[ab]

[a]*Department of Chemistry, University College London*
[b]*Thomas Young Centre, University College London*

*e05-discov-**dos***

Transparent conducting oxides (TCOs) combine optical transparency and electrical conductivity, making them indispensable in the fields of optoelectronics. However, the limited number of post-transition metal TCOs restricts the range of devices they can support. Recently, Ga-doped $ZnSb_2O_6$, an Sb(V)-based oxide with unique band alignment, has been computationally and experimentally examined, offering much-needed diversity to the field.[1] Inspired by this success, an underexplored post-transition binary Sb(V)-based system, Sb_2O_5 is considered a promising candidate due to its wide band gap and comparable electronic structures to the other successful TCOs.

In this work, we use hybrid density functional theory to investigate the electronic structure of Sb_2O_5, where wide fundamental and optical band gaps are found, enabling transparency. Furthermore, we apply the Phonopy to prove the dyn219(th)-5((h)-3(o)

A computational study characterising the intrinsic and extrinsic doping profile of ZnS

Savya

Organosulfide-Halide Perovskites incorporating Zwitterions

Santanu Saha[1,*], Jiayi Li[2], Hemamala I. Karunadasa[2,3], Marina R. Filip[1]

[1]*Department of Physics, University of Oxford, Clarendon Laboratory, Oxford OX1 3PU, United Kingdom*
[2]*Department of Chemistry, Stanford University, Stanford, CA 94305, United States*
[3]*Stanford Institute for Materials and Energy Sciences, SLAC National*

Prediction of critical micelle concentrations in surfactant mixtures by DPD simulations

Guadalupe Jiménez-Serratos[1,*], Annalaura del Regno[1], Patrick B. Warren[1], David J. Bray[1], Scott Singleton[2], and Richard L. Anderson[1]

1

Iron-sulfur peptides for an efficient electron transfer to the hydrogenase enzyme

Umberto Terranova*

*

How bulk and surface properties of Ti_4SiC_3, V_4SiC_3, Nb_4SiC_3 and Zr_4SiC_3 tune reactivity: A computational study

Matthew G. Quesne*, Nora H. de Leeuw, C. Richard A. Catlow

* *School of Chemistry, Cardiff University* *e05-react-**cat***

The -phase of over 100 ternary carbides/nitrides was first identified in the 1960s.[1] They display a very novel combination of properties including high conductivity and extreme resistance to oxidation or heating, which has led to these materials being lab [2] Structurally, the surfaces of these materials strongly resemble the pristine (111) facets of early transition metal carbides (TMCs), which have themselves been shown to be efficient catalysts for the hydrogenation of CO_2. Our group has previously undertaken a detailed in silico study into the catalytic activity of the equivalent TMCs to the MAX-phase material under investigation in this work and found the metal terminated (111) facets to be extremely active for CO_2 reduction.[3] The choice of the transition metal components was partly informed by this study and partly by a systematic screening of the bulk and surface properties of a diverse array of carbides.[4] This talk will focus on results published as part of a recent Faraday Discussion will present the results from a computational study into the bulk and catalytic properties of Ti_4SiC_3, V_4SiC_3, Nb_4SiC_3 & Zr_4SiC_3 and compare these properties to those already obtained for TiC, VC, NbC & ZrC.[5] We will show that the addition of an intestinal silicon layer into the bulk of these material increases the lattice paraments beyond those observe for the corresponding carbides. We also show that the pristine surfaces of the MAX-phase materials are much more active towards CO_2, H_2, H_2O and OH. However, our results demonstrate that pristine surfaces are not likely to be present in an oxygenating environment. Instead, the modelling predicts all surfaces will oxidise and barriers for CO_2 reduction will increase in the presence of H_2O, O_2 or OH. Current, work supported by the MCC is looking into the activity of these oxidized surfaces.

References

[1] W. Jeitschko, H. Nowotny and F. Benesovsky, *Monatshefte für Chemie*, 1963, **94**, 672–676.
[2] Z. M. Sun, H. Hashimoto, Z. F. Zhang, S. L. Yang and S. Tada, *Mater. Trans*., 2006, **47**, 170–174.

Posters

Advances in sustainable catalysis: A computational perspective

Matthew G. Quesne*, Fabrizio Silveri, Nora H. de Leeuw, C. Richard A. Catlow

* *School of Chemistry, Cardiff University* *e05-react-**cat***

The urgent need for society to move towards a greener and more sustainable future presents a very exciting opportunity for catalytic chemists. Many of the necessary changes in resource management and increased energy efficiency will be propelled by the directed design of new catalysts, for which a detailed theoretical understanding of

Capturing the ground state electronic and magnetic properties of Cobalt containing metal and mixed metal oxides using density functional theory (DFT).

Akash Hiregange*, Pavel Stishenko, Andrew Logsdail, James Paterson, Sarah Haigh

**School of Chemistry, Cardiff University.*

*e05-surfin-**log***

The Fischer-

Modelling hole trapping in hydrogen-related defects at the c-Si/a-SiO_2 and c-Si/a-SiO_2/a-HfO_2 interface

Teofilo Cobos Freire and Alexander Shluger

*Department of Physics and Astronomy, University College London e05-surfin-shl

The performance metal-oxide-semiconductor field-effect transistors (MOSFETs) is affected by random telegraph noise and bias temperature instability. These effects are caused by the trapping and release of holes and electrons in the dielectric oxide layer,

Computational Infrared and Raman Spectroscopy in ChemShell

Catalytic Partial Oxidation of Methane using an Yttria-Stabilized Zirconia Catalyst

Potential thermoelectric materials: zinc-based oxychalcogenides $SrZn_2Ch_2O$ (Ch = S, Se)

Shipeng Bi*, Katarina Brlec, David O. Scanlon

*Department of Chemistry, University College London *e05-power-**dos***

In the process of primary energy consumption, a large amount of waste heat is generated. Thermoelectric materials can convert waste heat energy into electric energy, which is crucial to solving the current energy crisis and global warming. In general, the performance of thermoelectric materials can be determined by the dimensionless figure of merit, ZT, that is evaluated by electronic and thermal properties. However, it is difficult to maximize ZT due to the interaction between different parameters. Although some high ZT thermoelectric materials such as Bi_2Te_3 and PbTe have been found, they have not been widely used because they contain rare or toxic elements.[1,2]

Recently, $SrZn_2S_2O$ was proposed as a novel photocatalyst for water splitting applications.[3] Because of its complex structure, relatively high thermal stability, and the low toxicity and easy synthesis of oxides, we think that $SrZn_2S_2O$ may be a potential thermoelectric material. Using density functional theory (DFT), we first studied the phononic stability of $SrZn_2S_2O$ and its variant - $SrZn_2Se_2O$. Then based on the $SrZn_2Ch_2O$ (Ch = S, Se) electronic structure calculated with hybrid-DFT, electronic transport properties were calculated using AMSET[4]. Our calculation results indicate that both oxychalcogenides may become thermoelectric materials for high-temperature applications.

References

1 Y. Z. Pei, A. D. LaLonde, N. A. Heinz and G. J. Snyder, *Adv. Energy Mater.*, 2012, **2**, 670-675.

2 M. Saleemi, M. Toprak, S. Li, M. Johnsson and M. Muhammed, *J. Mater. Chem.*, 2012, **22**, 725-730.

3 S. Nishioka, T. Kanazawa, K. Shibata, Y. Tsujimoto, H.-C. Loye and K. Maeda, *Dalton T.*, 2019, **48**, 15778–15781.

4 A. M. Ganose, J. Park, A. Faghaninia, R. Woods-Robinson, K. A. Persson and A. Jain, *Nat. Commun.*, 2021, **12**, 2222.

Influence of Solvent on Selective Catalytic Reduction of Nitrogen Oxides with Ammonia over Cu-CHA Zeolite

Jamal Abdul Nasir*, Alexey A. Sokol, Jingcheng Guan, You Lu, Thomas W. Keal, Andrew M. Beale, C. Richard A. Catlow

*Department of Chemistry, University College London e05-react-sok

In this new study, we made a comparison between water- and ammonia-solvated and bare Cu species [1]. Our results show the promoting effect of solvent on the oxidation component of the NH_3-SCR cycle since the formation of important nitrate species is found to be energetically more favourable on the solvated Cu. Conversely, both solvent molecules are predicted to inhibit the reduction component of the NH_3-SCR cycle. Diffuse reflectance infrared fourier-transform spectroscopy (DRIFTS) experiments exploiting (concentration) modulation excitation spectroscopy (MES) and phase-sensitive detection (PSD) identified spectroscopic signatures of Cu-nitrate and Cu-nitrosamine (H_2NNO), important species which had not been previously observed experimentally. This is further supported by the QM/MM-calculated harmonic vibrational analysis. Additional insights are provided into the reactivity of solvated active sites and the formation of key intermediates including their formation energies and vibrational

Removal of VOCs in cooking fumes via adsorption method on zeolites

Yuanyuan Zhang*, Alexey A. Sokols, C. Richard A. Catlow

*Department of Chemistry, University College London e05-enviro-cat

The growth of public awareness of hazard from gaseous pollutants emitted by cooking activities spawned the implement of new laws and regulations as well as new technologies for cooking fumes emission control. Considering the emission characteristics of large air volume and intermittence of cooking fumes, MFI zeolites were applied as an efficient adsorbent for enriching the low-concentrated complex humid volatile organic compounds (VOCs) from cooking fumes. In order to illustrate the superiority of Ti-O-Si framework as a cooking VOCs adsorbent, zeolites with the same MFI topology to TS-1 but different framework compositions of Si-O-Al (ZSM-5) and Si-O-Si (silicalite-1, hereinafter referred as S-1) were also investigated for comparison. Related characterizations were performed to reveal the difference among three zeolites. It was found that textural properties, surface silanol density and defect content significantly affect the adsorption site accessibility, water resistance as well as organophilicity of zeolites adsorbents, all of which play a decisive role in the adsorption process of cooking fumes. Furthermore, adsorption energy between water/VOCs molecular and zeolites with Si/Al/Ti center was calculated through hybrid quantum mechanical/molecular mechanical (QM/MM) technique, revealing the adsorption interaction between different framework and

Thermoelectric properties of the *Pnma* and *R3m* phases of GeS and GeSe

Min. Zhang*, Joseph M. Flitcroft, Jonathan M. Skelton

**Department of Chemistry, University of Manchester*

Stability of $Ru@Sn_9$ Zintl cluster on a CeO_2 (111) surface and its catalytic activity in Water-Gas-Shift (WGS) reaction.

Sourav Mondal*, John E. McGrady

*Department of Chemistry, University of Oxford e05-react-**mcg***

Cerium Oxide (CeO_2) is one of the most efficient compounds, supporting surface, because of the facile changes in oxidation state between Ce^{4+} to Ce^{3+}, and is widely used in high-performance oxygen storage applications and catalytic redox reactions, solid oxide fuel cells, water-gas shift reactions, etc. The use of Zintl clusters or metalloids rather than isolated noble metal atoms as the catalytic centre offers the potential to exploit the interactions between transition and main-group metal to achieve low-barrier reactions, and recent work from our collaborators in Tianjin suggests that Zintl clusters are effective catalysts for the water gas shift (WGS) reaction. Recently, Sun et al. have reported the selective reduction of CO_2 over highly dispersed $RuSnO_x$ sites, derived from a $[Ru@Sn_9]^{6-}$ Zintl cluster but an atomic-level understanding of mechanism, and how it relates to the electronic properties of the cluster remain unclear.[1]

In this work, we firstly investigate the stability of the $Ru@Sn_9$ Zintl cluster on the CeO_2 surface by comparing the optimised energy of the cluster on the surface with a dispersed structure where the Ru and Sn atoms are dispersed on the surface, using periodic Density Functional Theory (DFT). Interestingly, in many cases clusters prove not to be stable on surface, but rather are disrupted by strong metal-support interactions (SMSI).[2] An important first objective therefore is to establish the stability of the $Ru@Sn_9$ cluster on the surface under normal reaction conditions. It has been found that $Ru@Sn_9$ cluster will disperse over the surface and accumulate in a monolayer where Ru would prefer to stay at the edge. Later, we report some initial studies on the $RuSn_9/CeO_2$ (111) catalyst for WGS reaction where possible reaction mechanisms have been explored and activation energy barriers calculated. This catalyst shows a significantly lower energy barrier using different reaction path, compared to other reported heterogeneou[(int)ed heterogeneou

Utilisation of Machine Learning for Lithium Titanium Sulphate (LTS) Solid Solutions Energy Prediction

Stefani Alexandria Setiono, Bruno Camino, Scott M Woodley

[*]*Department of Chemistry, University*

Effects of Electronic Polarization on the Convergence of Mott-Littleton Calculations of $BaSnO_3$

Cyril Xu*, Alexey Sokol*, Scott Woodley*

*Department of Chemistry, University College London e05-bulk-**smw**

Barium Stannate ($BaSnO_3$) is a perovskite oxide material with a wide range of applications in various electronic devices thanks to its attractive electronic and thermal properties. Point defects in such materials can introduce additional charge carriers, modify band structures, and affect carrier mobility, resulting in changes in conductivities as well as catalytic activities. Understanding and controlling point defects in perovskite materials is crucial for tailoring their properties to specific applications and advancing their technological potential. The Mott-Littleton method evaluates the defect energy accurately by calculating the energy difference between the perfect and the defective crystals using interatomic potentials, which require significantly less computational resources, compared to *ab initio* methods. Partitioning the regions around the defect centre also provide straightforward and intuitive interpretations on defect energies based on factors that affect the formation and stability of the defect.

We show that converging the defect energy in Mott-Littleton calculations can be challenging due to effects of electronic polarization due to the charged point defect, even for simple crystal structures in bulk phase, and analysing charge distributions in localised cluster can provide further insights on how the model can be improved and adapted to surfaces and interfaces, where addressing defect-induced dipole and quadrupole effects remains problematic.

Ketonisation of Carboxylic Acids over ZrO_2-based catalysts for Biomass Valorisation: The Role of Surface Acid-Base Sites

M. Delarmelina*, G. Deshmukh, H. Manyar, C. R. A. Catlow

* *School of Chemistry, Cardiff University, Cardiff CF10 3AT, UK*

*e05-react-**cat***

Computational investigation of CO_2 hydrogenation over Rh-based catalysts

Shijia Sun*, Michael D. Higham, C. Richard A. Catlow

* *Department of Chemistry, University College London*

High-dimensional neural networks for interatomic potentials: Applications to metal organic frameworks

E. Kasoar*, D. Mason, P. Austin, A. Elena

**Scientific Computing, STFC UKRI*

Machine learning interatomic potentials (MLIP) are revolutionising our way of describing and understanding how atoms and molecules interact. Precise simulations of atomic and molecular chemical systems is hampered by the poor scalability of first principles

USING POTENTIALS FOR MODELLING COMPOUNDS THAT UNDERGO THE METAL – INSULATOR PHASE TRANSITION

Paula Gómez García

*Department of Chemistry, University College London e05-bulk-**smw***

The metal-insulator phase transition is the process by which a compound that conducts electricity becomes an insulator (or vice versa) by means of a change in its atomic and electronic structure. One of the most studied compounds is vanadium dioxide (VO_2). The computational method most widely used for its study is DFT since the process has its origin in a Peierls distortion caused by the presence of a d^1 electron in vanadium. Moreover, there are other dioxides that undergo the same process by means of a Peierls distortion: NbO_2 and TaO_2, being both niobium anm atin3(t)8(a)-3(til8(m)-6TJETQ

Modeling the Effect of OSDA on Nucleation Process of TS-1 Zeolite

Xu Zhang*; Sokols, Alexeykols

High-throughput computational design and discovery of conductive materials in the CSD MOF subset

Federica Zanca[1], Sanggyu Chong[2], Bartomeu Monserrat[3], David Fairen-Jimenez[4], Peyman Z. Moghadam[1*]

[1] Dept. of Chemical and Biological Engineering, University of Sheffield, Sheffield, United Kingdom;

e05-power

MOFs are innovative porous materials, and have been widely studied for the past two decades for applications in different areas including gas storage, gas separation and catalysis. Here, we aim to study electrical conductivity in MOFs, a less explored but interesting property that can bring promising opportunities in energy storage and sensing application. Due to their high porosity and surface area, MOFs are poor electrical conductors. Here, to identify promising conductive structures, we performed high-throughput screening of the existing ca. 90,000 structures in the CSD MOF subset[1] characterising the band gap and examining the presence or absence of metallic behavior. The first set of selection criteria was developed based on the nature secondary building unit and the ligand. We focused on MOFs containing open shell metals, metal clusters and highly conjugated linkers that can facilitate through-linker charge transfer between metals. The second set of criteria involved the presence of linkers containing metal-S, or -N coordination, redox- -
the ca. 1000 structures shortlisted, we then performed DFT calculations to derive useful insights into structure-conductivity relationships in MOFs, identify top-performing conductive MOFs, and to delineate key chemical and physical features in MOFs that influence their conductive properties for the first time. The results guide MOF researchers to assess and design conductive structures for electronics, energy storage and sensing applications.

Reference:

[1] Peyman Z. Moghadam *et al*. (2017) Chemistry of Materials, 29, 2618 2625

Aurora Cruz-Cabeza (acc)	
University of Durham	
Aurora Cruz-Cabeza	aurora.j.cruz-cabeza@durham.ac.uk
C. Richard A. Catlow	
Chemistry, UCL and Cardiff University	
R. Catlow Jamal Abdul Nasir (Thurs-Friday) Maicon Delarmelina Erze Gao Michael Higham Tony Kinoti Yunxin Li Matthew Quesne Shijia Sun Ruotao Yang Xu Zhang	c.r.a.catlow@ucl.ac.uk jamal.nasir.18@ucl.ac.uk delarmelinam@cardiff.ac.uk erze.gao.19@ucl.ac.uk m.highman@ucl.ac.uk t.kinoti@ucl.ac.uk yunxin.li.22@ucl.ac.uk quesnem@cardiff.ac.uk uccaunb@ucl.ac.uk ruotao.yang.22@ucl.ac.uk uccaxz1@ucl.ac.uk
Clotilde S. Cucinotta (clo)	
Department of Chemistry, Imperial College London	
Rashid Al-Heidous	rea19@ic.ac.uk
George Darling (dar)	
Department of Chemistry, University of Liverpool	
Ben Asher (Thurs-Friday) Andrij Vasylenko (Thurs-Friday)	benjamin.asher@liverpool.ac.uk andrij@liverpool.ac.uk
David O. Scanlon (dos)	
Department of Chemistry, University College London	
Savyasanchi Aggarwal Shipeng Bi Ke Li	uccagga@ucl.ac.uk shipeng.bi.22@ucl.ac.uk zccakli@ucl.ac.uk
Marina Filip (fil)	
Department of Physics, University of Oxford	
Santanu Saha (Thurs-Friday)	santanu.saha@physics.ox.ac.uk

Andrea Floris (flo)	
School of Maths and Physics, University of Lincoln	
Andrea Floris (Friday online only)	afloris@lincoln.ac.uk
Matthew Foulkes (fou)	
Imperial College	
Halvard Sutterud (Thurs-Friday)	h.sutterud21@imperial.ac.uk
John Harding (har)	
Department of Materials Science & Engineering, University of Sheffield	
John Harding (Thurs-Friday)	j.harding@sheffield.ac.uk
John Buckeridge (jbu)	
School of Engineering, London South Bank University	
John Buckeridge	j.buckeridge@lsbu.ac.uk
Kostya Trachenko (kos)	
Queen Mary University London	

School of Maths and Physics, University of Lincoln	
Matt Watkins (Wed-Thursday) Warren Lynch (Wed-Thursday)	mwatkins@lincoln.ac.uk wlynch@lincoln.ac.uk
John McGrady (mcg)	
Department of Chemistry, University of Oxford	
Sourav Mondal Yao Zhao	sourav.mondal@chem.ox.ac.uk yao.zhao@new.ox.ac.uk
Keith P. McKenna (mck)	
Department of Physics, University of York	
Alex Armstrong	alex.armstrong@york.ac.uk
Marco Molinari (mol)	
Department of Chemistry, University of Huddersfield	
M. Molinari (Thurs-Friday) Minh Khoa Ta	m.molinari@hud.ac.uk khoa.ta@hud.ac.uk
Erich Muller (mul)	
Imperial College	
Erich Muller	e.muller@imperial.ac.uk
Samuel Murphy (mur)	
University of Lancaster	
Ryan Kerr (Thursday)	r.kerr2@lancaster.ac.uk
Thomas Keal (COSEC)	

Natalia Martsinovich (nat)	
Department of Chemistry, University of Sheffield	
Natalia Martsinovich (Thurs-Friday)	n.martsinovich@sheffield.ac.uk
Nicolas Harrison (nic)	
Department of Chemistry, Imperial College	
Felix Mildner (Wed-Thursday)	f.mildner19@imperial.ac.uk
Nikolas Kaltsoyannis (nik)	
Department of Chemistry, University of Manchester	
Jiali Chen Ryan Demsey	jiali.chen@manchester.ac.uk ryan.dempsey@postgrad.manchester.ac.uk
Jonathan Skelton (ske)	
University of Manchester	
Jonathan Skelton Joseph Flitcroft Min Zhang	jonathan.skelton@manchester.ac.uk joseph.flitcroft@manchester.ac.uk min.zhang-19@postgrad.manchester.ac.uk
Alexander L. Shluger (shl)	
Department of Phy66.384 719hy66.383(al)4(t)anETdPh	

Sohayb Amin Paula Gomez-Garcia Yuhui Joley Lin Zirui Qiu Stefani Alexandria Setiono	sohayb.amin.22@ucl.ac.uk paula.garcia.22@ucl.ac.uk zccainc@ucl.ac.uk zirui.qiu.22@ucl.ac.uk uccaset@ucl.ac.uk
Alexey Sokol (sok)	
Department of Chemistry, University College London	
Alexey Sokol Alec Desmoutier Jingcheng Guan Xingfan Zhang Yuanyuan Zhang	a.sokol@ucl.ac.uk uccades@ucl.ac.uk jingcheng.guan@ucl.ac.uk xingfan.zhang.20@ucl.ac.uk uccazhc@ucl.ac.uk
Umberto Terranova (ter)	
University of Buckingham	
Umberto Terranova	umberto.terranova@buckingham.ac.uk
Lucy Whalley (luc)	
University of Northumbria	
Lucy Whalley (Wed-Thursday)	l.whalley@northumbria.ac.uk
David Willock (wil)	
University of Cardiff	
David Willock	willockd@cardiff.ac.uk
Invited Speakers	

Sergei Dudarev (UKAEA, UK)
Jon Wakelin (Thursday online)
Lupe Jiménez-Quero
Neil Allan