



To learn is to myelinate
Patrick Long and Gabriel Corfas
Science **346**, 298 (2014);
DOI: 10.1126/science.1261127

This copy is for your personal, non-commercial use only.

If you wish to distribute this article to others, you can order high-quality copies for your colleagues, clients, or customers by [clicking here](#).

Permission to republish or repurpose articles or portions of articles can be obtained by following the guidelines [here](#).

The following resources related to this article are available online at www.sciencemag.org (this information is current as of October 16, 2014):

Updated information and services, including high-resolution figures, can be found in the online version of this article at:

<http://www.sciencemag.org/content/346/6207/298.full.html>

A list of selected additional articles on the Science Web sites **related to this article** can be found at:

<http://www.sciencemag.org/content/346/6207/298.full.html#related>

This article **cites 12 articles**, 5 of which can be accessed free:

<http://www.sciencemag.org/content/346/6207/298.full.html#ref-list-1>

This article appears in the following **subject collections**:

Neuroscience

<http://www.sciencemag.org/cgi/collection/neuroscience>

Downloaded from www.sciencemag.org on October 16, 2014

gree to which axons are myelinated affect the timing of the flow of information between neurons. The composition of myelinated axons in the brain is heterogeneous; axons may be highly myelinated, not myelinated, or incompletely myelinated, and the pattern of myelination may vary along a single axon (4